

PCT


WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 401/06, 403/06, 413/06, 211/60, A61K 31/55, 31/445		A1	(11) International Publication Number: WO 96/31498
			(43) International Publication Date: 10 October 1996 (10.10.96)
(21) International Application Number: PCT/DK96/00139		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 1 April 1996 (01.04.96)			
(30) Priority Data: 0405/95 7 April 1995 (07.04.95) DK 1005/95 11 September 1995 (11.09.95) DK			
(71) Applicant: NOVO NORDISK A/S [DK/DK]; Novo Allé, DK-2880 Bagsværd (DK).			
(72) Inventors: DÖRWALD, Florenzio, Zaragossa; Klokkebyet 3D, DK-2730 Herlev (DK). ANDERSEN, Knud, Erik; Nøddehønden 122, DK-2765 Smørum (DK). MADSEN, Peter; Ulvejerg 7, DK-2880 Bagsværd (DK). JØRGENSEN, Tine, Krogh; Stavnshøj Alle 80, DK-2730 Herlev (DK). HOHLWEG, Rolf; Nybovej 6, DK-3490 Kvistgaard (DK). ANDERSEN, Henrik, Sune; Kastelsvej 24 st. th., DK-2100 København Ø (DK). TREPPENDAHL, Svend; Frederiksdalsvej 221, DK-2830 Virum (DK). OLSEN, Uffe, Bang; Hørsbred 111, DK-2625 Vallensbæk (DK). ZDENĚK, Polivka; Zborovská 23, 150 00 Praha 5 (CZ). ALEXANDRA, Silhánková; V Sthlách 1311/3, 142 000 Praha 4 (CZ). KAREL, Sindelár; Olbrachtova 50, 140 00 Praha 4 (CZ).		Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	

(54) Title: NOVEL HETEROCYCLIC COMPOUNDS



(57) Abstract

A compound of formula (I) in which Z is selected from (a), (b), (c), (d), (e), (f), (g), (h), (i), (j), and (k). The present invention relates to novel N-substituted azaheterocyclic carboxylic acids and esters thereof in which a substituted alkyl chain forms part of the N-substituent or salts thereof, to methods for their preparation, to compositions containing them, and to their use for the clinical treatment of painful, hyperalgesic and/or inflammatory conditions in which C-fibers play a pathophysiological role by eliciting neurogenic pain or inflammation.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TC	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

Novel heterocyclic Compounds

Field of the Invention

The present invention relates to novel N-substituted azaheterocyclic carboxylic
5 acids and esters thereof in which a substituted alkyl chain forms part of the N-
substituent or salts thereof, to methods for their preparation, to compositions
containing them, and to their use for the clinical treatment of painful,
hyperalgesic and/or inflammatory conditions in which C-fibers play a
pathophysiological role by eliciting neurogenic pain or inflammation. The
10 invention also relates to the use of the present compounds for the treatment
of insulin resistance in non-insulin-dependent diabetes mellitus (NIDDM) or
aging, the present compounds knowing to interfere with neuropeptide con-
taining C-fibres and hence inhibit the secretion and circulation of insulin an-
tagonizing peptides like CGRP or amylin.

15 Background of the Invention

The nervous system exerts a profound effect on the inflammatory response.
Antidromic stimulation of sensory nerves results in localized vasodilation and
increased vascular permeability (Janecso et al. Br. J. Pharmacol. 1967, 31,
138-151) and a similar response is observed following injection of peptides
20 known to be present in sensory nerves. From this and other data it is postu-
lated that peptides released from sensory nerve endings mediate many inflam-
matory responses in tissues like skin, joint, urinary tract, eye, meninges,
gastro-intestinal and respiratory tracts. Hence inhibition of sensory nerve
peptide release and/or activity, may be useful in treatment of, for example
25 arthritis, dermatitis, rhinitis, asthma, cystitis, gingivitis, thrombo-phlebitis,

Further, the potent effects of CGRP on skeletal muscle glycogen synthase activity and muscle glucose metabolism, together with the notion that this peptide is released from the neuromuscular junction by nerve excitation, suggest that CGRP may play a physiological role in skeletal muscle glucose metabolism by directing the phosphorylated glucose away from glycogen storage and into the glycolytic and oxidative pathways (Rossetti et al. *Am. J. Physiol.* **264**, E1-E10, 1993). This peptide may represent an important physiological modulator of intracellular glucose trafficking in physiological conditions, such as exercise, and may also contribute to the decreased insulin action and skeletal muscle glycogen synthase activity in pathophysiological conditions like NIDDM or aging-associated obesity (Melnyk et al. *Obesity Res.* **3**, 337-344, 1995) where circulating plasma levels of CGRP are markedly increased. Hence inhibition of release and/or activity of the neuropeptide CGRP may be useful in the treatment of insulin resistance related to type 2 diabetes or aging.


In US Patent No. 4,383,999 and No. 4,514,414 and in EP 236342 as well as in EP 231996 some derivatives of N-(4,4-disubstituted-3-butenyl)azaheterocyclic carboxylic acids are claimed as inhibitors of GABA uptake. In EP 342635 and EP 374801, N-substituted azaheterocyclic carboxylic acids in which an oxime ether group and vinyl ether group forms part of the N-substituent respectively are claimed as inhibitors of GABA uptake. Further, in WO 9107389 and WO 9220658, N-substituted azacyclic carboxylic acids are claimed as GABA uptake inhibitors. EP 221572 claims that 1-aryloxyalkylpyridine-3-carboxylic acids are inhibitors of GABA uptake.

In addition to the above cited references, US Patent No. 3,074,953 discloses 1-(3-(10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-1-propyl)-4-phenyl-4-piperidinecarboxylic acid ethyl ester as a psychotropic drug. Analogous 1-substituted 4-phenyl-4-piperidinecarboxylic acid ester derivatives to the above cited compound are described (*J. Med. Chem.* 1967, 10, 627-635 and *J. Org. Chem.* 1962, 27, 230-240) as analgesics, antispasmodics and psychotropics. In JP

49032544, JP 48040357, FR 2121423, GB 1294550 and DE 2101066, 1-substituted 4-dialkylamino-4-piperidinecarboxamides are disclosed as psychotropic agents, for the treatment of schizophrenia and as inhibitors of inflammation.

Description of the Invention

The present invention relates to novel N-substituted azaheterocyclic carboxylic acids and esters thereof of formula I



5 wherein R¹ and R² independently are hydrogen, halogen, trifluoromethyl, NR⁶R⁷, hydroxy, C₁-₆-alkyl or C₁-₆-alkoxy; and

Y is >N-CH₂-, >CH-CH₂- or >C=CH- wherein only the underscored atom participates in the ring system; and


X is -O-, -S-, -C(R⁶R⁷)-, -CH₂CH₂-, -CH=CH-CH₂-, -CH₂-CH=CH-, -CH₂-(C=O)-,

10 -(C=O)-CH₂-, -CH₂CH₂CH₂-, -CH=CH-, -N(R⁸)-(C=O)-, -(C=O)-N(R⁸)-, -O-CH₂-, -CH₂-O-, -S-CH₂-, -CH₂-S-, -(C=O)-, -N(R⁹)- or -(S=O)- wherein R⁶, R⁷, R⁸ and R⁹ independently are hydrogen or C₁-₆-alkyl; and

r is 1, 2 or 3; and

Z is selected from

- 4 -



wherein n is 1 or 2; and

R^3 is $-(CH_2)_mOH$ or $-(CH_2)_pCOR^4$ wherein m is 0, 1, 2, 3, 4, 5 or 6 and p is 0 or 1 and wherein R^4 is -OH, -NH₂, -NHOH or C₁₋₆-alkoxy; and

5 R^5 is hydrogen, halogen, trifluoromethyl, hydroxy, C₁₋₆-alkyl or C₁₋₆-alkoxy; and

R^{10} is hydrogen, C₁₋₆-alkyl, C₁₋₆-alkoxy or phenyl optionally substituted with halogen, trifluoromethyl, hydroxy, C₁₋₆-alkyl or C₁₋₆-alkoxy; and

R^{11} is hydrogen or C₁₋₆-alkyl; and

... is optionally a single bond or a double bond;

10 or a pharmaceutically acceptable salt thereof.

- 5 -

The compounds of formula I may exist as geometric and optical isomers and all isomers and mixtures thereof are included herein. Isomers may be separated by means of standard methods such as chromatographic techniques or fractional crystallization of suitable salts.

5

Preferably, the compounds of formula I exist as the individual geometric or optical isomers.

The compounds according to the invention may optionally exist as pharmaceutically acceptable acid addition salts or - when the carboxylic acid group is not esterified - as pharmaceutically acceptable metal salts or - optionally alkylated - ammonium salts.

Examples of such salts include inorganic and organic acid addition salts such as hydrochloride, hydrobromide, sulphate, phosphate, acetate, fumarate, maleate, citrate, lactate, tartrate, oxalate or similar pharmaceutically acceptable inorganic or organic acid addition salts, and include the pharmaceutically acceptable salts listed in Journal of Pharmaceutical Science, 66, 2 (1977) which are hereby incorporated by reference.

20

The term "C₁₋₆-alkyl" as used herein, alone or in combination, refers to a straight or branched, saturated hydrocarbon chain having 1 to 6 carbon atoms such as e.g. methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, 2-methylbutyl, 3-methylbutyl, n-hexyl, 4-methylpentyl, neopentyl, n-hexyl, 1,2-dimethylpropyl, 2,2-dimethylpropyl and 1,2,2-trimethylpropyl.

25

- 6 -

The term "C₁₋₆-alkoxy" as used herein, alone or in combination, refers to a straight or branched monovalent substituent comprising a C₁₋₆-alkyl group linked through an ether oxygen having its free valence bond from the ether oxygen and having 1 to 6 carbon atoms e.g. methoxy, ethoxy, propoxy, isopropoxy,
5 butoxy, pentoxy.

The term "halogen" means fluorine, chlorine, bromine or iodine.

Illustrative examples of compounds encompassed by the present invention
10 include:

1-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-3-piperidinecarbox-
amide;

15 1-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-4-piperidinecarboxylic
acid;

1-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-2-piperidinecarboxylic
acid;

20 (1-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-3-piperidinyl)methanol;

4-(4-Chlorophenyl)-1-(3-(10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-4-pi-
peridinol;

25

- 7 -

4-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-2-piperazinecarboxylic acid;

(2S,4R)-1-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-4-hydroxy-2-
5 pyrrolidinecarboxylic acid;

4-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-2-morpholinecarboxylic acid;

10 1-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-2-aziridinecarboxylic acid;

2-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-1,2,3,4-tetrahydro-4-isoquinolinecarboxylic acid;

15

1-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-4-methyl-[1,4]-diazepane-6-carboxylic acid;

2-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-1,2,3,4-tetrahydro-3-
20 isoquinolinecarboxylic acid;

1-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-3-piperidinecarboxylic acid hydroxamide;

25 (4-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)piperazin-1-yl)acetic acid;

- 8 -

1-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-4-piperidineacetic acid;

1-(3-(10,11-Dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-1-propyl)-4-piperidinecarboxylic acid;

5

(R)-1-(3-(10,11-Dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-1-propyl)-3-piperidinecarboxamide;

(R)-1-(3-(10,11-Dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-1-propyl)-2-pyrrolidinecarboxylic acid;

10

(S)-1-(3-(10,11-Dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-1-propyl)-2-pyrrolidinecarboxylic acid;

15 (R)-1-(3-(10,11-Dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-1-propyl)-2-piperidinecarboxylic acid;

1-(3-(10H-Phenoxazin-10-yl)-1-propyl)-4-piperidinecarboxylic acid;

20 1-(3-(3-Chloro-10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-4-piperidinecarboxylic acid;

1-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-3-piperidineacetic acid;

25

- 9 -

1-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-2-methyl-3-piperidinecarboxylic acid;

1-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-3-quinuclidinium-
5 carboxylate;

1-(3-(2,8-Dibromo-10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-4-piperidinecarboxylic acid;

10 1-(3-(3,7-Dichloro-10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-4-piperidinecarboxylic acid;

1-(3-(3-Methyl-10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-4-piperidinecarboxylic acid;

15

1-(3-(3,7-Dimethyl-10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-4-piperidinecarboxylic acid;

1-(3-(3-Dimethylamino-10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-4-
20 piperidinecarboxylic acid;

or a pharmaceutically acceptable salt thereof.

As used herein, the term "patient" includes any mammal which could benefit
25 from treatment of neurogenic pain or inflammation or insulin resistance in
NIDDM. The term particularly refers to a human patient, but is not intended to
be so limited.

- 10 -

- It has been demonstrated that the novel compounds of formula I inhibit neurogenic inflammation which involves the release of neuropeptides from peripheral and central endings of sensory C-fibres. Experimentally this can be demonstrated in animal models of formalin induced pain or paw oedema
- 5 (Wheeler and Cowan, Agents Actions 1991, 34, 264-269) in which the novel compounds of formula I exhibit a potent inhibitory effect. Compounds of formula I may be used to treat all painful, hyperalgesic and/or inflammatory conditions in which C-fibers play a pathophysiological role by eliciting neurogenic pain or inflammation, i.e.:
- 10 Acutely painful conditions exemplified by migraine, postoperative pain, burns, bruises, post-herpetic pain (Zoster) and pain as it is generally associated with acute inflammation; chronic, painful and/or inflammatory conditions exemplified by various types of neuropathy (diabetic, post-traumatic, toxic), neuralgia, rheumatoid arthritis, spondylitis, gout, inflammatory bowel disease, prostatitis,
- 15 cancer pain, chronic headache, coughing, asthma, chronic pancreatitis, inflammatory skin disease including psoriasis and autoimmune dermatoses, osteoporotic pain.

- Further, it has been demonstrated that the compounds of general formula I improves the glucose tolerance in diabetic ob/ob mice and that this may result from the reduced release of CGRP from peripheral nervous endings. Hence the compounds of general formula I may be used in the treatment of NIDDM as well as aging-associated obesity. Experimentally this has been demonstrated by the subcutaneous administration of glucose into ob/ob mice with or without
- 25 previous oral treatment with a compound of general formula I.

The compounds of formula I may be prepared by the following method:

- 11 -



- A compound of formula II wherein R^1 , R^2 , X, Y and r are as defined above and W is a suitable leaving group such as halogen, p-toluene sulphonate or mesylate may be reacted with an azaheterocyclic compound of formula III
- 5 wherein Z is as defined above. This alkylation reaction may be carried out in a solvent such as acetone, dibutylether, 2-butanone, methyl ethyl ketone, ethyl acetate, tetrahydrofuran (THF) or toluene in the presence of a base e.g. sodium hydride and a catalyst, e.g. an alkali metal iodide at a temperature up to reflux temperature for the solvent used for e.g. 1 to 120 h. If esters have been
- 10 prepared in which R^4 is alkoxy, compounds of formula I wherein R^4 is OH may be prepared by hydrolysis of the ester group, preferably at room temperature in a mixture of an aqueous alkali metal hydroxide solution and an alcohol such as methanol or ethanol, for example, for about 0.5 to 6 h.
- 15 Compounds of formula II and III may readily be prepared by methods familiar to those skilled in the art.

Under certain circumstances it may be necessary to protect the intermediates

- 12 -

used in the above methods e.g. a compound of formula III with suitable protecting groups. The carboxylic acid group can, for example, be esterified. Introduction and removal of such groups is described in "Protective Groups in Organic Chemistry" J.F.W. McOmie ed. (New York, 1973).

5

Pharmacological Methods

Formalin induced pain or paw oedema

Values for *in vivo* inhibition of formalin induced pain or oedema for the compounds of the present invention were assessed in mice essentially by the method of Wheeler-Aceto and Cowan (Agents Action 1991, 34, 265-269).

About 20 g NMRI female mice were injected 20 µl 1 % formalin into the left hind paw. The animals were then placed on a heated (31 °C) table, and the pain response was scored. After 1 h they were killed and bled. Left and right hind paws were removed and the weight difference between the paws was used as indication of the oedema response of the formalin injected paw.

Reduced release of CGRP

20

ob/ob female mice, 16 weeks of age, were injected glucose (2g/kg) subcutaneously. At times hereafter blood glucose was determined in tail venous blood by the glucose oxidase method. At the end of the study the animals were decapitated and trunk blood collected. Immunoreactive CGRP was determined in plasma by radio-immuno-assay. Two groups of animals were used. The one group was vehicle treated, whereas the other group received a compound of formula I via drinking water (100 mg/l) for five days before the test.

- 13 -

Values for inhibition of formalin induced pain response for some representative compounds are recorded in table 1.

TABLE 1

5 Inhibition of formalin induced pain response at 0.1 mg/kg

	Example no.	% Pain inhibition
	2	13
	4	47
10	5	36
	8	34
	9	29

- 15 For the above indications the dosage will vary depending on the compound of formula I employed, on the mode of administration and on the therapy desired. However, in general, satisfactory results are obtained with a dosage of from about 0.5 mg to about 1000 mg, preferably from about 1 mg to about 500 mg of compounds of formula I, conveniently given from 1 to 5 times daily, optional-
- 20 ly in sustained release form. Usually, dosage forms suitable for oral administration comprise from about 0.5 mg to about 1000 mg, preferably from about 1 mg to about 500 mg of the compounds of formula I admixed with a pharmaceutical carrier or diluent.
- 25 The compounds of formula I may be administered in a pharmaceutically acceptable acid addition salt form or where possible as a metal or a lower alkylammo-

- 14 -

nium salt. Such salt forms exhibit approximately the same order of activity as the free base forms.

. This invention also relates to pharmaceutical compositions comprising a compound of formula I or a pharmaceutically acceptable salt thereof and, usually, such compositions also contain a pharmaceutical carrier or diluent. The compositions containing the compounds of this invention may be prepared by conventional techniques and appear in conventional forms, for example capsules, tablets, solutions or suspensions.

10

The pharmaceutical carrier employed may be a conventional solid or liquid carrier. Examples of solid carriers are lactose, terra alba, sucrose, talc, gelatin, agar, pectin, acacia, magnesium stearate and stearic acid. Examples of liquid carriers are syrup, peanut oil, olive oil and water.

15

Similarly, the carrier or diluent may include any time delay material known to the art, such as glyceryl monostearate or glyceryl distearate, alone or mixed with a wax.

20 If a solid carrier for oral administration is used, the preparation can be tableted, placed in a hard gelatin capsule in powder or pellet form or it can be in the form of a troche or lozenge. The amount of solid carrier will vary widely but will usually be from about 25 mg to about 1 g. If a liquid carrier is used, the preparation may be in the form of a syrup, emulsion, soft gelatin capsule or
25 sterile injectable liquid such as an aqueous or non-aqueous liquid suspension or solution.

Generally, the compounds of this invention are dispensed in unit dosage form

- 15 -

comprising 50-200 mg of active ingredient in or together with a pharmaceutically acceptable carrier per unit dosage.

The dosage of the compounds according to this invention is 1-500 mg/day, e.g.

- 5 about 100 mg per dose, when administered to patients, e.g. humans, as a drug.

A typical tablet which may be prepared by conventional tableting techniques contains

10

Core:

Active compound (as free compound or salt thereof)	100 mg
Colloidal silicon dioxide (Areosil®)	1.5 mg
15 Cellulose, microcryst. (Avicel®)	70 mg
Modified cellulose gum (Ac-Di-Sol®)	7.5 mg
Magnesium stearate	

Coating:

20 HPMC	approx.	9 mg
Mywacett® 9-40 T	approx.	0.9 mg

*Acylated monoglyceride used as plasticizer for film coating.

- 25 The route of administration may be any route which effectively transports the active compound to the appropriate or desired site of action, such as oral or parenteral e.g. rectal, transdermal, subcutaneous, intranasal, intramuscular, topical, intravenous, intraurethral, ophthalmic solution or an ointment, the oral

- 16 -

route being preferred.

EXAMPLES

- 5 The process for preparing compounds of formula I and preparations containing them is further illustrated in the following examples, which, however, are not to be construed as limiting.

Hereinafter, TLC is thin layer chromatography, CDCl_3 is deuterio chloroform
10 and DMSO-d_6 is hexadeuterio dimethylsulfoxide. The structures of the compounds are confirmed by either elemental analysis or NMR, where peaks assigned to characteristic protons in the title compounds are presented where appropriate. ^1H NMR shifts (δ_{H}) are given in parts per million (ppm). M.p. is melting point and is given in $^{\circ}\text{C}$ and is not corrected. Column chromatography
15 was carried out using the technique described by W.C. Still et al, J. Org. Chem. (1978), 43, 2923-2925 on Merck silica gel 60 (Art. 9385). Compounds used as starting materials are either known compounds or compounds which can readily be prepared by methods known per se.


EXAMPLE 1

20

1-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-4-piperidinecarboxylic acid hydrochloride

25

- 17 -



- 5 To a suspension of 10,11-dihydro-5H-dibenz[b,f]azepine (15.2 g, 0.078 mol) in toluene (100 ml), 3-chloropropionyl chloride (9.50 ml, 0.099 mol) was added, and the resulting mixture was heated at reflux temperature for 1 h. Saturated aqueous sodium bicarbonate (100 ml) was added, and the phases were separated. The organic phase was washed with brine (100 ml), dried (MgSO_4) and concentrated in vacuo. This afforded 23.6 g of 3-chloro-1-(10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propanone as a solid which was used in the following step without further purification.

M.p. 107-108 °C.

- 15 Calculated for $\text{C}_{17}\text{H}_{16}\text{ClNO}$:
C, 71.45 %; H, 5.64 %; N, 4.90 %. Found:
C, 71.45 %; H, 5.79 %; N, 5.01 %.

- To a solution of 3-chloro-1-(10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propanone (14.0 g, 0.044 mol) in tetrahydrofuran (150 ml) at 0 °C, sodium borohydride (6.66 g, 0.176 mol) was added, followed by dropwise addition of glacial acetic acid (10.0 ml). The resulting mixture was stirred at room tempera-
- 20

- 18 -

- ture overnight and then heated at reflux temperature for 2 h. More sodium borohydride (6.50 g, 172 mmol) and then borontrifluoride diethyl etherate (20.0 ml, 0.163 mol) were added and heating at reflux temperature was continued for 20 h. Water (350 ml) was cautiously added and the phases were separated.
- 5 The aqueous phase was extracted with toluene (3 x 100 ml). The combined organic phases were washed with brine (3 x 100 ml), dried (MgSO_4) and concentrated in vacuo. The residue was purified by flash chromatography on silica gel (100 g) using a gradient of heptane and ethyl acetate (10:0 \rightarrow 10:2), to give 4.58 g (38 %) of 5-(3-chloropropyl)-10,11-dihydro-5H-dibenz[b,f]azepine
- 10 as an oil.

TLC: R_f = 0.63 (SiO_2 ; ethyl acetate/heptane = 1:2).

- A mixture of 4-piperidinecarboxylic acid ethyl ester (2.55 g, 16.2 mmol),
- 15 acetonitrile (13 ml), the above chloride (2.00 g, 0.0074 mol) and potassium iodide (1.14 g, 0.0068 mol) was heated at reflux temperature for 4 h and then stirred at room temperature overnight. Water (50 ml) was added and the product was extracted with ethyl acetate (3 x 20 ml). The combined organic extracts were washed with brine (2 x 20 ml), dried (MgSO_4) and concentrated
- 20 in vacuo. The residue was purified by flash chromatography on silica gel using a gradient of heptane and ethyl acetate (10:1 \rightarrow 1:1), to give 1.6 g (54 %) of 1-(3-(10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-4-piperidinecarboxylic acid ethyl ester as an oil.

- 25 TLC: R_f = 0.26 (SiO_2 ; ethyl acetate/heptane = 1:1).

Calculated for $\text{C}_{25}\text{H}_{32}\text{N}_2\text{O}_2$:

C, 76.50 %; H, 8.22 %; N, 7.14 %. Found:

C, 76.34 %; H, 8.51 %; N, 6.88 %.

- 19 -

The above ester (1.01 g, 2.57 mmol) was dissolved in ethanol (10 ml), and a solution of sodium hydroxide (0.59 g, 14.8 mmol) in water (1.5 ml) was added. The resulting mixture was stirred at room temperature for 3.5 h. A mixture of water (20 ml) and concentrated hydrochloric acid (3.0 ml) was added, and the
5 aqueous phase was extracted with dichloromethane (3 x 15 ml). The combined organic extracts were washed with brine (20 ml) and dried (MgSO₄). Evaporation of the solvent gave a foam, which was redissolved in a mixture of methanol (1.0 ml) and ethyl acetate (5.0 ml). Concentration in vacuo afforded a solid, which was suspended in ethyl acetate (15 ml), heated at reflux tempera-
10 ture for 1 minute and allowed to cool to room temperature. The solid was filtered off and dried to give 0.9 g (88 %) of the title compound as a powder.

Mp 195-197 °C.

Calculated. for C₂₃H₂₈N₂O₂.HCl:


15 C, 68.90 %; H, 7.29 %; N, 6.99 %. Found:

C, 68.90 %; H, 7.55 %; N, 6.72 %.

EXAMPLE 2

20

4-(4-Chlorophenyl)-1-(3-(10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)piperidin-4-ol hydrochloride



Ethyl malonyl chloride (25.0 g, 0.166 mol) was added to a suspension of 10,11-
5 dihydro-5H-dibenz[b,f]azepine (27.6 g, 0.141 mol) in toluene (250 ml), and the
resulting mixture was heated at reflux temperature for 1 h. Saturated, aqueous
sodium bicarbonate (200 ml) was added, and the phases were separated. The
organic phase was washed with brine (2 x 150 ml), dried (MgSO_4) and concen-
trated in vacuo. This afforded 56.0 g (100 %) of 3-(10,11-dihydro-5H-
10 dibenz[b,f]azepin-5-yl)-3-oxopropionic acid ethyl ester as an oil, which was used
in the following step without further purification.

Lithium aluminum hydride (20.0 g, 0.527 mol) was introduced into a
roundbottom flask under a nitrogen atmosphere. Toluene (800 ml) was added,
15 followed by tetrahydrofuran (80 ml). The resulting suspension was cooled to 10-
20 °C. The above crude 3-oxopropionic acid ester was dissolved in
tetrahydrofuran (250 ml) and slowly added dropwise. The addition rate was
adjusted to assure that the temperature was kept at 10-20 °C. The resulting
mixture was stirred at room temperature overnight. After cooling, 2 N sodium
20 hydroxide (200 ml) was added cautiously. Water (1.0 l) was added, the organic
layer was decanted off and the aqueous phase was extracted with toluene (2 x
300 ml). The combined organic phases were washed with brine (2 x 100 ml),
dried (MgSO_4) and concentrated in vacuo. The residue was purified by flash
chromatography on silica gel (175 g) using a gradient of heptane and ethyl
25 acetate (10:0 → 2:1), affording 21.2 g (59 %) of 3-(10,11-dihydro-5H-dibenz-
[b,f]azepin-5-yl)-1-propanol as an oil.

To a stirred solution of the above alcohol (1.01 g, 0.004 mol) and triethylamine

- 21 -

(1.02 g, 0.010 mol) in toluene (25 ml) at 0°C, methanesulfonyl chloride (0.6 ml, 0.0077 mol) was added dropwise over 10 minutes. The resulting mixture was stirred at 0 °C for 1.5 h. Water (50 ml) was added and the phases were separated. The aqueous phase was extracted with toluene (50 ml), and the
5 combined organic phases were washed with brine (2 x 50 ml), dried (MgSO₄) and concentrated in vacuo. The crude mesylate was mixed with 4-(4-chlorophenyl)piperidin-4-ol (0.81 g, 0.004 mol) and potassium carbonate (1.08 g, 0.008 mol) in acetonitrile (9 ml) and heated at reflux temperature for 6 h. The reaction mixture was left stirring at room temperature for 2 days. Water (50 ml)
10 was added and the mixture was extracted with ethyl acetate (3 x 15 ml), washed with brine (2 x 20 ml), dried (MgSO₄) and concentrated in vacuo. Water (50 ml) was added and the mixture was acidified by addition of concentrated hydrochloric acid (3 ml). The aqueous solution was extracted with dichloromethane (2 x 20 ml), washed with brine (2 x 20 ml), dried (MgSO₄) and
15 concentrated in vacuo. The resulting oil was dissolved in a mixture of ethyl acetate (15 ml) and methanol (2 ml). Heptane was added in small portions, just until the solution turned slightly turbid. After 4 h, crystals were filtered off, washed with heptane and dried, affording 1.1 g (61 %) of the title compound as a solid.

20

M.p. 189-191 °C.

Calculated for C₂₈H₃₀N₂O, HCl:

C, 69.56 %; H, 6.67 %; N, 5.79 %; Found:

C, 69.88 %; H, 6.92 %; N, 5.62 %.


25

EXAMPLE 3

1-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-3-piperidinemethanol

- 22 -

hydrochloride



5

A mixture of 3-(hydroxymethyl)piperidine (1.01 g, 0.0088 mol), acetonitrile (9 ml), 5-(3-chloropropyl)-10,11-dihydro-5H-dibenzo[b,f]azepine (0.86 g, 0.003 mol, pre-prepared similarly as described in example 1) and potassium iodide (0.56 g, 0.003 mol) was heated at reflux temperature for 18 h. Water (20 ml) was added, and the mixture was extracted with ethyl acetate (3 x 15 ml). The combined organic extracts were washed with brine (2 x 20 ml), dried (MgSO₄) and concentrated in vacuo. The residue was redissolved in a mixture of water (20 ml) and concentrated hydrochloric acid (3 ml), and extracted with dichloro-
15 methane (3 x 10 ml). The combined organic extracts were washed with brine (20 ml), dried (MgSO₄) and concentrated in vacuo. The residue was crystallised from a mixture of methanol (0.5 ml) and ethyl acetate (5 ml), affording 0.8 g (61 %) of the title compound as needles.

20 M.p. 145-147 °C.

Calculated for C₂₃H₃₀N₂O, HCl:

C, 71.39; H, 8.07; N, 7.24; Found:


- 23 -

C, 71.15; H, 8.29; N, 7.01

EXAMPLE 4

5

1-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-3-piperidine-carboxamide hydrochloride



10

To a stirred solution of 3-(10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propanol (1.44 g, 0.0057 mol, prepared similarly as described in example 2) and triethylamine (1.46 g, 0.014 mol) in toluene (40 ml) at 0°C, methanesulfonyl chloride (0.88 ml, 0.011 mol) was added dropwise over 10 minutes. The resulting mixture was stirred at 0 °C for 1.5 h. Toluene (50 ml) and water (100 ml) were added, and the phases were separated. The aqueous phase was extracted with toluene (50 ml), and the combined organic phases were washed with brine (2 x 100 ml), dried (MgSO₄) and concentrated in vacuo. The residue was dissolved in acetonitrile (20 ml) and 3-piperidinecarboxamide (1.09 g, 0.0085 mol) and potassium carbonate (1.76 g, 0.013 mol) were added. The

- 24 -

mixture was heated at reflux temperature for 4 h, and stirred at room temperature for 40 h. Water (20 ml) was added, and the product was extracted with ethyl acetate (2 x 20 ml). The combined organic extracts were washed with brine (2 x 20 ml), dried (MgSO_4) and concentrated in vacuo. Water (20 ml) and concentrated hydrochloric acid (3.0 ml) were added, and the mixture was extracted with dichloromethane (2 x 20 ml), dried (MgSO_4) and concentrated in vacuo. The residue was redissolved in a mixture of warm ethyl acetate (10 ml) and methanol (1.0 ml), and after standing for 5 h at room temperature, a precipitate was filtered off and dried, affording 1.56 g (64 %) of the title compound.


HPLC retention time = 20.44 minutes (5 μm C18 4 x 250 mm column, eluting with a 20-80 % gradient of 0.1 % trifluoroacetic acid/acetonitrile and 0.1 % trifluoroacetic acid/water over 30 minutes at 35 $^{\circ}\text{C}$).

Calculated for $\text{C}_{23}\text{H}_{29}\text{N}_3\text{O}$, HCl, 1.5 H_2O :
C, 64.70 %; H, 7.78 %; N, 9.84 %; Found:
C, 65.13 %; H, 7.85 %; N, 9.85 %.

EXAMPLE 5

1-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-2-piperidinecarboxylic acid hydrochloride

- 25 -



- 5 A mixture of 2-piperidinecarboxylic acid ethyl ester hydrochloride (0.60 g, 0.003 mol), acetonitrile (10 ml), 5-(3-chloropropyl)-10,11-dihydro-5H-dibenzo[b,f]azepine (0.60 g, 0.002 mol, prepared similarly as described in example 1), potassium iodide (0.40 g, 0.002 mol), potassium carbonate (1.03 g, 0.008 mol) and N,N-dimethylformamide (5 ml) was heated at reflux temperature
- 10 for 85 h. Water (50 ml) was added, and the aqueous solution was extracted with ethyl acetate (3 x 20 ml). The combined organic extracts were washed with brine (2 x 50 ml), dried (MgSO₄) and concentrated in vacuo. The product was purified by column chromatography on silica gel (15 g) using a gradient of heptane and ethyl acetate (100:0 --> 100:25), affording 0.83 g (97 %) of 1-(3-
- 15 (10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-2-piperidinecarboxylic acid ethyl ester as an oil.

TLC: R_f = 0.58 (SiO₂: heptane/ethyl acetate = 1:1).

- 20 The above ester (0.83 g, 0.002 mol) was dissolved in a mixture of ethanol (8 ml), water (2 ml) and sodium hydroxide (0.58 g, 0.015 mol). The reaction mixture was stirred at room temperature for 50 h and at 50 °C for 5 h. Water

- 26 -

(50 ml) and concentrated hydrochloric acid (3 ml) were added, and the resulting mixture was extracted with dichloromethane (3 x 10 ml). The combined organic extracts were washed with brine (2 x 20 ml), dried (MgSO_4) and concentrated in vacuo. The residue was re-dissolved in methanol (2 ml) and ethyl acetate (5 ml) and concentrated in vacuo. The solid residue was washed with a small amount of ethyl acetate and dried, affording 0.6 g (73 %) of the title compound as a powder.


M.p. 122-126 °C.

- 10 Calculated for $\text{C}_{23}\text{H}_{28}\text{N}_2\text{O}_2$, HCl, 0.25 H_2O :
C, 68.14 %; H, 7.33 %; N, 6.91 %; Found:
C, 68.34 %; H, 7.63 %; N, 6.66 %.

EXAMPLE 6

15

Potassium 4-(3-(10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-2-piperazine-carboxylate



- 27 -

A mixture of 2-piperazinecarboxylic acid dihydrochloride (5.06 g, 0.025 mol), ethanol (100 ml) and concentrated sulphuric acid (6.0 ml) was heated at reflux temperature for 6 days. Toluene (10 ml) was added, and the resulting mixture was concentrated in vacuo to 2/3 of its original volume. Cold, saturated aqueous potassium carbonate (80 ml) was added and the mixture was extracted with toluene (3 x 100 ml). The combined organic extracts were washed with brine (30 ml), dried (MgSO₄) and concentrated in vacuo, affording 1.0 g (26 %) of 2-piperazinecarboxylic acid ethyl ester as an oil. The oil crystallised upon standing at room temperature.

10

To a stirred solution of 3-(10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propanol (1.17 g, 0.0046 mol, prepared similarly as described in example 2) and triethylamine (1.18 g, 0.012 mol) in toluene (30 ml) at 0°C, methanesulfonyl chloride (0.70 ml, 0.009 mol) was added dropwise over 10 minutes. The resulting mixture was stirred at 0 °C for 1 h. Toluene (50 ml) and water (100 ml) were added, and the phases were separated. The aqueous layer was extracted with toluene (50 ml), and the combined organic phases were washed with brine (2 x 100 ml), dried (MgSO₄) and concentrated in vacuo. The residue was dissolved in acetonitrile (10 ml) and the above 2-piperazine-carboxylic acid ethyl ester (1.40 g, 0.0089 mol), potassium carbonate (0.67 g, 0.0049 mol) and toluene (5 ml) were added. The resulting mixture was heated at reflux temperature for 18 h. Water (50 ml) was added, and the mixture was extracted with ethyl acetate (3 x 20 ml). The combined organic extracts were washed with brine (2 x 20 ml), dried (MgSO₄) and concentrated in vacuo. The residue was purified by column chromatography on silica gel (27 g) using a gradient of methanol and ethyl acetate (5:100 → 20:100), affording 0.6 g (33 %) of 4-(3-(10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-2-piperazinecarboxylic acid ethyl ester as an oil.

25

- 28 -

TLC: $R_f = 0.60$ (SiO_2 ; ethyl acetate/methanol = 1:1).

A mixture of the above ester (0.55 g, 0.0014 mol), ethanol (5 ml), water (1 ml) and sodium hydroxide (0.34 g, 0.0085 mol) was stirred at room temperature for 5 18 h. Water (50 ml) and concentrated hydrochloric acid (3 ml) were added and the solution was washed with dichloromethane (4 x 15 ml). The dichloromethane extracts were discarded. The aqueous phase was made alkaline by addition of potassium carbonate (7.1 g) and the mixture was extracted with dichloromethane (4 x 15 ml). The combined organic extracts 10 were washed with brine (20 ml), dried (MgSO_4) and concentrated *in vacuo*. This gave 0.6 g of an oil, which was triturated with acetonitrile (2 ml) and then dried *in vacuo*, affording 0.5 g (90 %) of the title compound as a waxy solid.

M.p. 151-155 °C.

15 Calculated for $\text{C}_{22}\text{H}_{26}\text{N}_3\text{O}_3\text{K}$, 0.5 H_2O :


C, 64.05 %; H, 6.59 %; N, 10.18 %; Found:

C, 64.34 %; H, 7.04 %; N, 10.16 %.

EXAMPLE 7

20

Sodium 4-(3-(10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-1-piperazineacetate



- 29 -

To a solution of 3-(10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propanol (1.44 g, 0.0057 mol, prepared similarly as described in example 2) in dichloromethane (30 ml) at 0 °C, triethylamine (1.73 g, 0.017 mol) was added followed by methanesulfonyl chloride (0.9 ml, 0.012 mol). The resulting mixture was stirred for 30 minutes at 0 °C. Water (50 ml) was added, the phases were separated, and the organic layer was washed with brine (20 ml), dried (MgSO₄) and concentrated in vacuo. The residue was dissolved in acetonitrile (10 ml), N-(ethoxycarbonylmethyl)piperazine (2.44 g, 0.014 mol) was added and the reaction mixture was heated at 82 °C for 5.5 h. Water (50 ml) was added and the mixture was extracted with ethyl acetate (3 x 20 ml). The combined organic extracts were washed with brine (30 ml), dried (MgSO₄) and concentrated in vacuo. The residue was purified by column chromatography on silica gel (35 g) using ethyl acetate as eluent. This gave 1.5 g (63 %) of 4-(3-(10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-1-piperazineacetic acid ethyl ester as an oil.

TLC: R_f = 0.18 (SiO₂: ethyl acetate).

20

A mixture of the above ester (0.94 g, 0.0023 mol), ethanol (5 ml), water (1 ml) and sodium hydroxide (0.31 g, 0.0078 mol) was stirred at room temperature for 4.5 h. Water (50 ml) and concentrated hydrochloric acid (3 ml) were added and the solution was washed with dichloromethane (3 x 10 ml). The organic extracts were discarded. The aqueous phase was made alkaline by addition of 4 N sodium hydroxide (20 ml) and potassium carbonate (4 g). The mixture was extracted with dichloromethane (5 x 20 ml), and the combined organic extracts were dried (MgSO₄) and concentrated in vacuo, affording 0.9 g of a foam. The

- 30 -

foam was triturated and washed with ethyl acetate (2 ml), and dried in vacuo, affording 0.5 g (56 %) of the title compound as a solid.


HPLC retention time = 17.81 minutes (5 μ m C18 4 x 250 mm column, eluting with a 20-80 % gradient of 0.1 % trifluoroacetic acid/acetonitrile and 0.1 % trifluoroacetic acid/water over 30 minutes at 35 °C)

¹H NMR (400 MHz, CDCl₃) δ 1.60 (m, 2H), 2.10 - 2.60 (m, 12H), 3.12 (bs, 4H), 3.68 (m, 2H), 6.83 - 7.17 (m, 8H).

10

EXAMPLE 8

4-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-2-morpholinecarboxylic acid hydrochloride



15

A mixture of 2-morpholinecarboxylic acid ethyl ester (0.50 g, 0.0031 mol, prepared similarly as described in Tetrahedron Letters, Volume 32, 2281-4, 1991), acetonitrile (6 ml), potassium carbonate (0.50 g, 0.0036 mol), potassium iodide (0.54 g, 0.0033 mol), methanesulfonic acid 3-(10,11-dihydro-5H-

20

- 31 -

dibenz[b,f]azepin-5-yl)-1-propyl ester (0.36 g, 0.0011 mol, prepared similarly as described in example 2) and 5-(3-chloropropyl)-10,11-dihydro-5H-dibenz[b,f]azepine (0.40 g, 0.0015 mol, prepared similarly as described in example 1) was heated at reflux temperature for 22 h. Water (50 ml) was added, and the mixture was extracted with ethyl acetate (3 x 15 ml). The combined organic phases were washed with brine (20 ml), dried (MgSO_4) and concentrated in vacuo. The product was purified by column chromatography on silica gel (30 g) using a gradient of heptane and ethyl acetate (10:0 \rightarrow 10:4). This gave 0.4 g (42 %) of 4-(3-(10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-2-morpholinecarboxylic acid ethyl ester as an oil.

TLC: R_f = 0.33 (SiO_2 : ethyl acetate/heptane = 1:1).

A mixture of the above ester (0.40 g, 0.0010 mol), ethanol (10 ml) and 4 N sodium hydroxide (2 ml) was stirred at room temperature for 17 h. Water (50 ml) and concentrated hydrochloric acid (3 ml) were added and the mixture was extracted with ethyl acetate (3 x 20 ml). The combined organic extracts were washed with brine (20 ml) and dried (MgSO_4). After filtering off the drying agent, a solid started to precipitate from the solution. After standing for 4 h at room temperature, the precipitate was filtered off, and the product was dried in vacuo. This gave 0.2 g (49 %) of the title compound as a solid.

M.p. 196-199 °C.


Calculated for $\text{C}_{22}\text{H}_{26}\text{N}_2\text{O}_3$, HCl, 0.25 H_2O :

C, 64.86 %; H, 6.80 %; N, 6.88 %; Found:
C, 65.12 %; H, 7.09 %; N, 6.39 %.

EXAMPLE 9

- 32 -

2-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-1,2,3,4-tetrahydro-3-isoquinolinecarboxylic acid hydrochloride



5

To a solution of 3-(10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propanol (1.05 g, 0.0041 mol, prepared similarly as described in example 2) in dichloromethane (40 ml) at 0 °C, triethylamine (1.28 g, 0.013 mol) was added, followed by methanesulfonyl chloride (0.9 ml, 0.012 mol). After stirring for 30 minutes at 0 °C, water (50 ml) was added and the phases were separated. The organic layer was washed with brine (20 ml), dried (MgSO₄) and concentrated in vacuo. The residue was dissolved in acetonitrile (12 ml), and 1,2,3,4-tetrahydro-3-isoquinolinecarboxylic acid methyl ester (1.13 g, 0.0050 mol), N,N-dimethylformamide (5 ml), potassium carbonate (1.32 g, 0.0096 mol) and potassium iodide (0.30 g, 0.0018 mol) were added. The reaction mixture was heated at 82 °C for 12 h. N,N-Dimethylformamide (5 ml) was added and heating was continued for further 16 h. Water (50 ml) was added and the mixture was extracted with ethyl acetate (3 x 20 ml). The combined organic extracts were washed with brine (2 x 20 ml), dried (MgSO₄) and concentrated in vacuo. The product was purified by column chromatography on silica gel (80 g) using a

- 33 -

gradient of heptane and ethyl acetate (10:0 --> 10:3), affording 1.4 g (76 %) of 2-(3-(10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-1,2,3,4-tetrahydro-3-isoquinolinecarboxylic acid methyl ester as an oil.

5 TLC: R_f = 0.61 (SiO₂: heptane/ethyl acetate = 1:1).

A mixture of the above ester (0.80 g, 0.0019 mol), ethanol (5 ml), tetrahydrofuran (5 ml) and 4 N sodium hydroxide (4 ml) was stirred at room temperature for 22 h. Water (50 ml) and concentrated hydrochloric acid (2 ml) were added and the mixture was extracted with ethyl acetate (2 x 20 ml). The combined organic extracts were washed with brine (10 ml), dried (MgSO₄) and concentrated in vacuo. This gave 0.8 g of a solid, which was triturated and washed with ethyl acetate (2 x 5 ml). Drying in vacuo, afforded 0.6 g (75 %) of the title compound as a solid.

15

M.p. 205-208 °C.

Calculated for C₂₇H₂₈N₂O₂, HCl, 0.25 H₂O:

C, 71.51 %; H, 6.56 %; N, 6.18 %; Found:

C, 71.34 %; H, 6.69 %; N, 5.90 %.


20

EXAMPLE 10

1-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-4-piperidineacetic acid hydrochloride

25

- 34 -



A mixture of 5-(3-chloropropyl)-10,11-dihydro-5H-dibenz[b,f]azepine (1.5 g,
5 0.0055 mol, prepared similarly as described in example 1) and potassium
iodide (5.4 g, 0.0327 mol) in methyl ethyl ketone (100 ml) was heated at reflux
temperature for 2.5 h. Potassium carbonate (1.5 g, 0.0109 mol) and
4-piperidineacetic acid ethyl ester (1.4 g, 0.0082 mmol, described in J. Am.
Chem. Soc., Vol. 75, 6249, 1953) were added and the reaction mixture was
10 stirred at 75 °C overnight. After cooling, the reaction mixture was filtered (Hyflo)
and the solvent was evaporated in vacuo. The residue was purified by flash
chromatography on silica gel (150 g) using a gradient of heptane and ethyl
acetate (1:1→ 3:7), to give 0.6 g of 1-(3-(10,11-dihydro-5H-dibenz[b,f]-
azepin-5-yl)-1-propyl)-4-piperidineacetic acid ethyl ester as an oil.
15
TLC: R_f = 0.10 (SiO_2 : ethyl acetate/heptane = 1:1).

To a solution of the above ester (0.6 g, 0.0015 mol) in ethanol (5 ml) 4 N
sodium hydroxide (0.8 ml) was added and the mixture was stirred at room
20 temperature for 2.5 h and then left in a freezer overnight. The cold reaction
mixture was allowed to warm to room temperature during 1 h, and 4 N hydro-
chloric acid (1.2 ml) and water (10 ml) were added. The mixture was xtracted

- 35 -

with dichloromethane (2 x 100 ml). The combined organic extracts were dried (MgSO₄) and the solvent was evaporated in vacuo. The residue was reevaporated with a mixture of acetone and isopropyl acetate and then treated with a mixture of acetone and isopropyl acetate to give a solid which was
5 isolated and dried in vacuo. This afforded 0.33 g of the title compound.

M.p. 185-188 °C.


Calculated. for C₂₄H₃₀N₂O₂·HCl:

C, 69.47 %; H, 7.53 %; N, 6.75 %. Found:

10 C, 69.21 %; H, 7.80 %; N, 6.45 %.

EXAMPLE 11

1-(3-(10,11-Dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-1-propyl)-4-
15 piperidinecarboxylic acid hydrochloride



20

5-(3-Bromo-1-propylidene)-10,11-dihydro-5H-dibenzo[a,d]cycloheptene (3.00 g,

- 36 -

0.0096 mol), potassium carbonate (8.3 g, 0.060 mol), potassium iodide (3.3 g, 0.020 mol) and 4-piperidinecarboxylic acid ethyl ester (3.1 ml, 0.020 mol) was mixed in methyl ethyl ketone (100 ml) and heated at reflux temperature for 20 h and stirred at room temperature for 3 days. Water (100 ml) was added, the
5 phases were separated, and the aqueous phase was extracted with ethyl acetate. The organic phase was dried (MgSO_4) and evaporated in vacuo to give the crude product in almost quantitative yield. The crude product was treated with 1 N hydrochloric acid and ethyl acetate, evaporated to dryness and crystallised from ethyl acetate to give 2.89 g of 1-(3-(10,11-dihydro-5H-
10 dibenzo[a,d]cyclohepten-5-ylidene)-1-propyl)-4-piperidinecarboxylic acid ethyl ester hydrochloride as a solid.

M.p. 169-170 ° C.

15 A mixture of the above ester (0.105 g, 0.27 mmol), ethanol (15 ml) and 1 N sodium hydroxide (10 ml) was heated at reflux temperature for 3 h and then cooled to room temperature. Water (75 ml) was added, and the mixture was acidified with 5 N hydrochloric acid and extracted with dichloromethane (3 x 75 ml), dried (MgSO_4) and evaporated in vacuo. The resulting foam was
20 crystallised from acetone affording 0.080 g of the title compound as crystals.

M.p. 224 - 226° C.

Calculated. for $\text{C}_{24}\text{H}_{27}\text{NO}_2\cdot\text{HCl}$:


C, 72.44 %; H, 7.09 %; N, 3.52 %. Found:

25 C, 72.83 %; H, 7.38 %; N, 3.23 %.

EXAMPLE 12

- 37 -

(R)-1-(3-(10,11-Dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-1-propyl)-3-piperidinecarboxamide hydrochloride



5

(R)-1-(3-(10,11-Dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-1-propyl)-3-piperidinecarboxylic acid hydrochloride (4.96 g, 12.5 mmol, prepared as described in WO 9518793) was dissolved in N,N-dimethylformamide (60 ml). N-Hydroxybenzotriazole (1.86 g, 13.8 mmol) and 1-ethyl-3-(3-dimethylamino-propyl)carbodiimide hydrochloride (2.64 g, 13.8 mmol) were added, and the resulting mixture was stirred at room temperature for 20 minutes. Ammonium hydrogencarbonate (1.98 g, 25 mmol) was added and the mixture was stirred 1 day at room temperature. Ethyl acetate (200 ml) was added and the resulting mixture was extracted with water (200 ml), 5 % aqueous citric acid (200 ml), and saturated sodium hydrogencarbonate (200 ml). The combined aqueous phases were evaporated to dryness in vacuo and the residue was extracted with dichloromethane (200 ml). The resulting dichloromethane suspension was filtered and evaporated. The residue was purified by column chromatography on silica gel (600 ml) using a mixture of ethyl acetate and triethyl amine (95:5) as eluent. This afforded 1.92 g (43 %) of the free base as an oil, which was

- 38 -

converted to the hydrochloride by dissolution in diethyl ether (25 ml) and addition of 1 N hydrochloric acid in diethyl ether (5.9 ml). Filtration followed by drying in vacuo, afforded 1.53 g, (31 %) of the title compound as a solid.

- 5 TLC: R_f = 0.33 (SiO₂: Ethyl acetate/triethylamine = 95:5).
¹H NMR (400 MHz, CDCl₃): δ 1.38 (dq, 1H), 1.75 - 1.97 (m, 3H), 2.5 - 3.7 (m, 13H), 5.78 (t, 1H), 7.0 - 7.3 (m, 8H), 7.63 (s, 1H), 10.8 (s, 1H).


M.p. >250 °C

- 10 Calculated for C₂₄H₂₈N₂O, HCl:
C, 72.62 %; H, 7.36 %; N, 7.06 %; Found:
C, 72.24 %; H, 7.59 %; N, 6.87 %.

15

EXAMPLE 13

(R)-1-(3-(10,11-Dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-1-propyl)-2-pyrrolidinecarboxylic acid hydrochloride



20

- 39 -

5-(3-Bromo-1-propylidene)-10,11-dihydro-5H-dibenzo[a,d]cycloheptene (3.86 g, 12.3 mmol, prepared as described in WO 9518793), potassium carbonate (10.2 g, 74 mmol), potassium iodide (4.08 g, 24.6 mmol), and D-proline methyl ester hydrochloride (2.45 g, 14.8 mmol) were mixed in methyl ethyl ketone (40 ml) and heated at reflux temperature for 20 h. After cooling, the mixture was filtered and the filtrate was evaporated in vacuo. The residue was purified by column chromatography on silica gel (800 ml) using a mixture of ethyl acetate and heptane (1:3) as eluent. This afforded 0.96 g (22 %) of (R)-1-(3-(10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-1-propyl)-2-pyrrolidinecarboxylic acid methyl ester as an oil.

TLC: R_f = 0.39 (SiO₂: ethyl acetate/heptane = 1:2).

15

The above ester was dissolved in 1,4-dioxane (40 ml), and water (5 ml) was added. 1 N Aqueous sodium hydroxide (2.15 ml) was added in portions at room temperature over 6 h. The mixture was stirred overnight at room temperature. 1 N Aqueous sodium hydroxide (0.84 ml) was added in portions over 20 h. Water (100 ml) was added and the mixture was washed with diethyl ether (2 x 100 ml). The aqueous phase was acidified to pH = 2 with 1 N hydrochloric acid and extracted with dichloromethane (2 x 100 ml). The combined organic extracts were dried (MgSO₄) and evaporated in vacuo to give 0.29 g (41 %) of the title compound as an amorphous solid.

25

¹H-NMR (400 MHz, CDCl₃): δ 1.95 (bs, 2H), 2.3 (m, 2H), 2.5-3.4 (m, 11 H), 5.78 (t, 1H), 7.0-7.3 (m, 8H).

- 40 -

Calculated for $C_{23}H_{25}NO_2 \cdot HCl$:

C, 71.96 %; H, 6.83 %; N, 3.65 %, Found:


C, 72.15 %; H, 7.37 %; N, 3.40 %.

5

EXAMPLE 14

(S)-1-(3-(10,11-Dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-1-propyl)-2-pyrrolidinecarboxylic acid hydrochloride

10



-
- 15 5-(3-Bromo-1-propylidene)-10,11-dihydro-5H-dibenzo[a,d]cycloheptene (2.00 g, 6.4 mmol, prepared as described in WO 9518793), potassium carbonate (5.3 g, 38.4 mmol), potassium iodide (2.12 g, 12.8 mmol), and L-proline methyl ester hydrochloride (1.27 g, 7.7 mmol) were mixed in methyl ethyl ketone (40 ml) and heated at reflux temperature for 12 h. After cooling, the mixture was
- 20 filtered, and the filtrate was evaporated. The residue was purified by column chromatography on silica gel (800 ml) using a mixture of ethyl acetate and

- 41 -

heptane (1:2) as eluent. This afforded 1.64 g (71 %) of (S)-1-(3-(10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-1-propyl)-2-pyrrolidinecarboxylic acid methyl ester as an oil.

5 TLC: R_f = 0.39 (SiO₂: ethyl acetate/heptane = 1:2)

The above ester (1.3 g, 3.6 mmol) was dissolved in 1,4-dioxane (50 ml), and water (5 ml) was added. 1 N Aqueous sodium hydroxide (3.8 ml) was added in portions at room temperature over 6 h. The mixture was stirred overnight at
10 room temperature. Water (100 ml) was added and the mixture was washed with diethyl ether (2 x 100 ml). The aqueous phase was acidified to pH = 2 with 1 N hydrochloric acid and extracted with dichloromethane (2 x 100 ml). The combined organic extracts were dried (MgSO₄) and evaporated *in vacuo* to give 0.80 g (58 %) of the title compound as an amorphous solid.

15

¹H-NMR (400 MHz, CDCl₃): δ 1.95 (bs, 2H), 2.3 (m, 2H), 2.5-3.4 (m, 11 H), 5.78 (t, 1H), 7.0-7.3 (m, 8H).


Calculated for C₂₃H₂₅NO₂, HCl:

20 C, 71.96 %; H, 6.83 %; N, 3.65 %; Found:
C, 72.61 %; H, 7.30 %; N, 3.34 %.

EXAMPLE 15

25 1-(3-(10,11-Dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-1-propyl)-2-piperidinecarboxylic acid hydrochloride.

- 42 -



- 5 5-(3-Bromo-1-propylidene)-10,11-dihydro-5H-dibenzo[a,d]cycloheptene (10.0 g, 22 mmol, prepared as described in WO 9518793), potassium carbonate (18.24 g, 132 mmol), and DL-pipecolinic acid ethyl ester hydrochloride (5.40 g, 26 mmol) were mixed in ethyl acetate (50 ml) and heated at reflux temperature for 16 h. Additional potassium carbonate (10 g), DL-pipecolinic acid ethyl ester
- 10 hydrochloride (2 g), and ethyl acetate (50 ml) were added and the mixture was heated at reflux temperature for 24 h. After cooling, the mixture was filtered, and the filtrate was evaporated. The residue was purified by column chromatography on silica gel (500 ml), eluting first with dichloromethane, and then with ethyl acetate. This afforded 7.92 g (92 %) of 1-(3-(10,11-dihydro-5H-
- 15 dibenzo[a,d]cyclohepten-5-ylidene)-1-propyl)-2-piperidine carboxylic acid ethyl ester as an oil.

TLC: $R_f = 0.54$ (SiO_2 ; ethyl acetate/heptane = 3:7)

- 20 1 N Hydrochloric acid (100 ml) was added to the above ester (7.7 g, 20 mmol) and the mixture was heated for 1 h while water and ethanol was distilled off (20 ml distilled). Water (40 ml) was added and the mixture was heated for 5 h while

- 43 -

water and ethanol was distilled off. After cooling, toluene (100 ml) was added to the mixture. The precipitated crystals were isolated by filtration and washed with 1 N hydrochloric acid, affording the crude product (5.3 g). A portion of this (2.0 g) was further purified by dissolution in water (100 ml, 75° C) and addition of 37 % hydrochloric acid (6 ml). The mixture was allowed to cool to room temperature. Filtration, washing with 1 N hydrochloric acid and drying in vacuo afforded 1.72 g of the title compound as a solid.

M.p. 126-127 °C.

10 ¹H NMR (300 MHz, CDCl₃): δ 1.5-1.75 (m, 4H), 2.1 (m, 1H), 2.5-3.5 (m, 11H), 4.10 (bs, 1H), 5.80 (t, 1H), 7.05-7.23 (m, 8H).

Calculated for C₂₃H₂₆NO₂, HCl, 0.5 H₂O:


C, 70.13 %; H, 7.16 %; N, 3.56 %, Found:

15 C, 69.74 %; H, 7.46 %; N, 3.08 %.

EXAMPLE 16

1-(3-(10H-Phenoxazin-10-yl)-1-propyl)-4-piperidinecarboxylic acid hydrochloride.

20



- 44 -

Phenoxazine (10.0 g, 54.6 mmol) was dissolved in N,N-dimethylformamide (300 ml) under a nitrogen atmosphere. Sodium hydride (3.27 g, 81.9 mmol, 60 % dispersion in oil) was added in portions, and the resulting mixture was stirred
5 for 20 minutes at room temperature. Dropwise, 1-bromo-3-chloro-propane (21.48 g, 0.136 mmol) was added. The mixture was stirred overnight. Ammoniumchloride (5.5 g, 0.10 mol) was added over 4 minutes and stirring was continued for 30 minutes. The mixture was poured into water (800 ml) and extracted with dichloromethane (2 x 600 ml). The combined organic extracts
10 were dried (MgSO_4) and evaporated in vacuo. This afforded 16.0 g crude 10-(3-chloropropyl)-10H-phenoxazine.

The above crude chloride (5.09 g, 17.4 mmol) was dissolved in acetonitrile (100 ml) and potassium iodide (2.74 g, 16.5 mmol) was added. 4-Piperidine-
15 carboxylic acid ethyl ester (6.00 g, 38.2 mmol) was dissolved in acetonitrile (30 ml) and added. The resulting mixture was heated at reflux temperature for 24 h and left stirring at room temperature for 48 h. Water (100 ml) was added followed by ethyl acetate (100 ml). The aqueous phase was extracted with ethyl acetate (3 x 100 ml), and the combined organic phases were washed with brine
20 (2 x 100 ml) and dried (MgSO_4). After evaporation in vacuo, the residue was purified by column chromatography on silica gel (500 ml) using a mixture of heptane and ethyl acetate (1:1) as eluent. This afforded 5.18 g (77 %) of 1-(3-(10H-phenoxazin-10-yl)-1-propyl)-4-piperidinecarboxylic acid ethyl ester as an oil.

25

TLC: R_f = 0.2 (SiO_2 : ethyl acetate/heptane = 1:1)

The above ester (1.56 g, 4.04 mmol) was dissolved in a mixture of 96 %

- 45 -

ethanol (20 ml) and tetrahydrofuran (20 ml). A solution of sodium hydroxide (0.95 g) in water (3 ml) was added, and the reaction mixture was stirred at room temperature for 1.5 h. 0.1 M Hydrochloric acid (28 ml) was added and the mixture was extracted with dichloromethane (3 x 30 ml). The combined organic
5 extracts were washed with brine (30 ml), dried (MgSO_4) and evaporated in vacuo. Twice, acetone was added and the solution was evaporated in vacuo. After a third addition of acetone, precipitation started, and the mixture was left stirring 2 h. After filtration, the solid was re-suspended in acetone (25 ml) and left stirring overnight. The solid was filtered off, washed with acetone and dried.
10 This afforded 1.30 g (83 %) of the title compound as a solid.

M.p. 196-198 °C.


Calculated for $\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_3$, HCl:

C, 64.86 %; H, 6.48 %; N, 7.20 %, Found:

15 C, 64.82 %; H, 6.78 %; N, 6.77 %.

EXAMPLE 17

20 1-(3-(3-Chloro-10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-4-piperidinecarboxylic acid hydrochloride.



- 46 -

3-Chloro-10,11-dihydro-5H-dibenz[b,f]azepine (3.82 g, 16.6 mmol) was dissolved in toluene (20 ml). A solution of 3-chloropropionylchloride (2.53 g, 19.9 mmol) in toluene was added dropwise, and the resulting mixture was heated to 95 °C and stirred at that temperature for 30 minutes. The mixture was stirred overnight at room temperature. Further 3-chloropropionylchloride (2.53 g, 19.9 mmol) was added and the mixture was stirred at 95 °C for 1.5 h. After cooling, 0.2 M sodium hydroxide (10 ml) was added, and the phases were separated.

10 The organic phase was diluted with more toluene (50 ml), and washed with first 0.2 M sodium hydroxide (6 x 10 ml) and then with more 0.2 M sodium hydroxide (3 x 20 ml) until the aqueous phase was alkaline. The organic phase was washed with water (3 x 15 ml), brine (25 ml), and dried (MgSO₄). Evaporation in vacuo afforded 5.23 g (98 %) of crude 3-chloro-1-(3-chloro-10,11-dihydro-5H-

15 dibenz[b,f]azepin-5-yl)-1-propanone as an oil. This was further purified by addition of a mixture of heptane and ethyl acetate (1:1). This afforded 3.14 g (59 %) of the product as a solid.

A 1.0 M solution of lithiumaluminium hydride in tetrahydrofuran (18.7 ml, 18.7 mmol) was introduced into a 250 ml dry, three-necked, roundbottom flask under a nitrogen atmosphere. The solution was cooled on an icebath. Concentrated sulphuric acid (0.5 ml) was added dropwise, with caution, over 10 minutes. More dry tetrahydrofuran (20 ml) was added to compensate for evaporated solvent and the mixture was stirred for 15 minutes. Additional tetrahydrofuran

25 was added (20 ml) and the icebath was removed. The mixture was stirred for 75 minutes at room temperature. The above amide (3.0 g, 9.3 mmol) was dissolved in dry tetrahydrofuran (25 ml) and dropwise added over 20 minutes. The reaction mixture was stirred for 1 h. Water (0.7 ml) was added, followed

- 47 -

subsequently by 4 N sodium hydroxide (0.7 ml) and water (2.1 ml). Stirring was continued for 30 minutes. The mixture was filtered (hyflo) and evaporated in vacuo, affording 2.70 g (95 %) 3-chloro-5-(3-chloropropyl)-10,11-dihydro-5H-dibenz[b,f]azepine as an oil.

5

The above chloride (1.50 g, 4.91 mmol) was dissolved in acetonitrile (10 ml), and a solution of 4-piperidinecarboxylic acid ethyl ester (1.70 g, 10.8 mmol) in acetonitrile (4 ml) was added, followed by potassium iodide (0.76 g, 4.6 mmol). The resulting mixture was heated at reflux temperature for 24 h. Water (50 ml) was added followed by ethyl acetate (50 ml). The aqueous phase was extracted with ethyl acetate (3 x 30 ml), and the combined organic phases were washed with brine (2 x 25 ml) and dried (MgSO_4). After evaporation in vacuo, the residue was purified by column chromatography on silica gel (150 ml) using a mixture of heptane and ethyl acetate (1:1) as eluent. This afforded 1.78 g (85 %)

15 % of 1-(3-(3-chloro-10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-4-piperidinecarboxylic acid ethyl ester as an oil.

TLC: R_f = 0.21 (SiO_2 ; ethyl acetate/heptane = 1:1)

20 The above ester (1.70 g, 3.98 mmol) was dissolved in 99 % ethanol (20 ml). A solution of sodium hydroxide (0.92 g) in water (2.5 ml) was added, and the reaction mixture was stirred at room temperature for 1 h. 1 N Hydrochloric acid (23 ml) was added and the mixture was extracted with dichloromethane (3 x 25 ml). The combined organic extracts were washed with brine (25 ml) and water (20 ml), dried (MgSO_4) and evaporated in vacuo. Dichloromethane was added and the solution was re-evaporated. Isopropyl acetate was added to the resulting foam, and the precipitated solid was filtered off and dried. This

25 afford d 1.28 g (74 %) of the crude title compound. The product was

- 48 -

redissolved in isopropanol, the solution was decanted and evaporated in vacuo. Dichloromethane was added and the solution was evaporated in vacuo. Isopropyl acetate was added to the resulting foam, and the precipitated solid was filtered off and dried. This procedure was repeated once more.

5

MS(EI) 398 ($M^+ - HCl$, 18 %).


HPLC retention time = 24.14 minutes (5 μ m C18 4 x 250 mm column, eluting with a 20-80 % gradient of 0.1 % trifluoroacetic acid/acetonitrile and 0.1 % trifluoroacetic acid/water over 30 minutes at 35 °C)

10

EXAMPLE 18

1-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-3-piperidineacetic acid hydrochloride

15



A suspension of 3-piperidineacetic acid (4.5 g, 0.032 mol, described in J. Org. Chem., 28, 602, 1963) in a mixture of dry hydrogen chloride (excess) in ethanol was stirred at ambient temperature. When the solid was dissolved, the solution was stirred for 2 days. The solvent was evaporated in vacuo and the residue

- 49 -

was reevaporated with diethyl ether (25 ml) and then stirred with diethyl ether (35 ml) for 20 minutes. The solid was isolated by filtration and dried. This afforded 6.1 g of 3-piperidineacetic acid ethyl ester hydrochloride as a solid.

5 M.p. 111-113 °C.

A mixture of potassium iodide (19.2 g, 0.12 mol) and methyl ethyl ketone (180 ml) was heated at reflux temperature for 1 h. A solution of 5-(3-chloropropyl)-10,11-dihydro-5H-dibenz[b,f]azepine (5.2 g, 0.019 mol, prepared similarly as described in example 1) in methyl ethyl ketone (25 ml) was added and heating at reflux temperature was continued for 3 h. Potassium carbonate (9.3 g, 0.067 mol) and 3-piperidineacetic acid ethyl ester hydrochloride (5.6 g, 0.027 mol) were added and the reaction mixture was heated at reflux temperature for 2 h. The temperature was lowered to just below reflux and the mixture was left with stirring overnight. After cooling, the reaction mixture was filtered (Hyflo) and the solvent was evaporated in vacuo. The residue was purified by flash chromatography on silica gel (225 g) using a mixture of heptane and ethyl acetate (1:1) as eluent, to give 5.0 g of 1-(3-(10,11-dihydro-5H-dibenz[b,f]-azepin-5-yl)-1-propyl)-3-piperidine-acetic acid ethyl ester as an oil.

TLC: R_f = 0.19 (SiO_2 ; ethyl acetate/heptane = 1:1).

To a solution of the above ester (2.5 g, 0.0062 mol) in ethanol (10 ml), 4 N sodium hydroxide (2.3 ml) was added and the mixture was stirred at room temperature for 3 h. 4 N Hydrochloric acid (3.8 ml) and water (10 ml) were added. The mixture was extracted with dichloromethane (2 x 250 ml). The combined organic extracts were dried (MgSO_4) and the solvent was evaporated

- 50 -

in vacuo. The residue was reevaporated twice with acetone and stirred with acetone for a while. The solid was isolated by filtration and dried. This afforded 2.4 g of the title compound as a solid.

5 M.p. 233-235 °C.

Calculated. for $C_{24}H_{30}N_2O_2 \cdot HCl$:

C, 69.47 %; H, 7.53 %; N, 6.75 %. Found:


C, 69.59 %; H, 7.78 %; N, 6.50 %.

10

EXAMPLE 19

1-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-2-methyl-3-piperidinecarboxylic acid hydrochloride

15



2-Methylnicotinic acid methyl ester (4.0 g, 0.026 mol) was dissolved in 1 N
20 hydrochloric acid (30 ml) and 10 % palladium on carbon (0.8 g) was added.
The resulting mixture was hydrogenated at 200 psi for 10 days. The reaction
mixture was filtered and the solid was washed with dichloromethane (100 ml)

- 51 -

and water (50 ml). The combined filtrates were evaporated in vacuo to give a residue which was reevaporated with dichloromethane (2 x 30 ml). This afforded 5.1 g of crude 2-methyl-3-piperidinecarboxylic acid methyl ester hydrochloride which was used for further reaction without purification.

5

A mixture of potassium iodide (17.5 g, 0.11 mol) and methyl ethyl ketone (180 ml) was heated at reflux temperature for 1 h. A solution of 5-(3-chloropropyl)-10,11-dihydro-5H-dibenz[b,f]azepine (4.8 g, 0.019 mol, prepared similarly as described in example 1) in methyl ethyl ketone (25 ml) was added and heating at reflux temperature was continued for 2 h. Potassium carbonate (8.5 g, 0.061 mol) and 2-methyl-3-piperidinecarboxylic acid methyl ester hydrochloride (5.1 g, 0.026 mol) were added and the reaction mixture was heated just below reflux temperature for 64 h. After cooling, the reaction mixture was filtered (Hyflo) and the solvent was evaporated in vacuo. The residue was purified by flash chromatography on silica gel (275 g) using a mixture of heptane and ethyl acetate (1:1) as eluent, to give 3.4 g of 1-(3-(10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-2-methyl-3-piperidinecarboxylic acid methyl ester as an oil.

20 To a solution of the above ester (3.4 g, 0.0087 mol) in 96 % ethanol (15 ml), 4 N sodium hydroxide (4.4 ml) was added and the mixture was stirred at room temperature for 4 h and then left in a refrigerator overnight. Stirring was continued at room temperature for 5 h and 4 N hydrochloric acid (6 ml) was added. The solvent was evaporated and the residue was triturated with acetone (30 ml) for 10 minutes. The solid was isolated by filtration, washed with acetone and dried in vacuo. The solid was suspended in a mixture of water (35 ml) and dichloromethane (800 ml). A saturated sodium bicarbonate solution was added until pH 8 - 9 and when the solid had dissolved. The phases were separated

- 52 -

and the aqueous phase was extracted with dichloromethane (250 ml). The organic extracts were combined and the volume was reduced by evaporation. Excess concentrated hydrochloric acid was added and the mixture was evaporated to dryness. The residue was reevaporated twice with dichloromethane and
5 then triturated with acetone. The solid was isolated by filtration and dried in vacuo. This afforded 2.4 g of the title compound as a solid.

M.p. 169-170 °C.

Calculated. for $C_{24}H_{30}N_2O_2 \cdot HCl \cdot H_2O$:


10 C, 66.59 %; H, 7.63 %; N, 6.47 %. Found:

C, 66.64 %; H, 7.94 %; N, 6.23 %.

EXAMPLE 20

15


1-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-3-quinuclidinium-carboxylate



- 5-(3-Chloropropyl)-10,11-dihydro-5H-dibenzo[b,f]azepine (1.70 g, 6.25 mmol, prepared as described in example 1), and quinuclidine-3-carboxylic acid methyl ester (0.85 g, 5.0 mmol) were dissolved in 2-butanone (25 ml). Dry potassium carbonate (4.15 g, 30 mmol) and sodium iodide (0.75 g, 5 mmol) were added and the stirred mixture was heated at reflux temperature for 5 h. After cooling to room temperature, toluene (25 ml) and water (25 ml) were added. An oily precipitate was formed, which was separated from the solvents by decanting and dissolved in dichloromethane (30 ml). The solution was washed with water (2 x 30 ml), 1 N hydrochloric acid (2 x 30 ml), saturated sodium hydrogencarbonate solution (30 ml) and water (30 ml). The organic layer was dried over MgSO_4 . Evaporation in vacuo afforded 1.15 g 1-(3-(10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-3-methoxycarbonyl-quinuclidinium hydroxide. The crude product was directly used in the next step without further purification.
- 15 The above ester (1.05 g, 2.50 mmol) was dissolved in ethanol (12.5 ml). A 2 N solution of sodium hydroxide (4.1 ml, 8.25 mmol) was added and the resulting mixture was stirred at room temperature for 10 minutes. Water (10 ml) was added and ethanol was evaporated in vacuo. The remaining aqueous solution was washed with diethyl ether (2 x 30 ml) and extracted with 1-butanol (3 x 20 ml). The 20 butanol phases were washed with water (10 ml) and evaporated in vacuo. Stripping of the remainder with n-heptane (3 x 20 ml) afforded the title compound (0.71 g) as a foam.
- 25 Calculated for $\text{C}_{25}\text{H}_{30}\text{N}_2\text{O}_2 \cdot 3.5 \text{H}_2\text{O}$:
C, 66.20 %; H, 8.22 %; N, 6.18 %; Found:
C, 66.52 %; H, 8.03 %; N, 5.79 %.

EXAMPLE 21

1-(3-(2,8-Dibromo-10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-4-piperidinecarboxylic acid hydrochloride



10

2,8-Dibromo-10,11-dihydro-5H-dibenz[b,f]azepine (1.41 g, 4.0 mmol, prepared according to K. Smith et al., Tetrahedron, 48, 7479 (1992)), was dissolved in toluene (25 ml). Triethylamine (0.60 ml, 4.4 mmol) and 3-chloropropionyl chloride (0.50 ml, 5.2 mmol) were added to the stirred solution at room temperature. Stirring was continued at room temperature for 1 h and at reflux temperature for 2.5 h. The reaction mixture was cooled, filtered and evaporated in vacuo. The crude 3-chloro-1-(2,8-dibromo-10,11-dihydro-5H-dibenz[b,f]azepine-5-yl)propan-1-one was used in the next step without further purification.

A solution of lithium aluminium hydride (0.326 g, 8.56 mmol) in dry tetrahydrofuran (30 ml) was cooled in an ice bath and concentrated sulphuric acid (0.428 g, 4.28 mmol) was added dropwise. The solution was stirred at room temperature for 0.5 h. A solution of the above product (1.90 g, 4.28
5 mmol) was added dropwise and stirring was continued for 0.5 h. The reaction was then quenched by careful addition of ethyl acetate (5 ml) followed by water (0.8 ml). Filtration of the mixture and evaporation of the filtrate in vacuo afforded 1.51 g (82 %) 2,8-dibromo-5-(3-chloropropyl)-10,11-dihydro-5H-dibenz[b,f]azepine as a foam, which was used in the next step without further
10 purification.

The above chloride (1.5 g, 3.5 mmol) and ethyl isonipecotate (0.79 g, 5.0 mmol) was dissolved in 2-butanone (40 ml). Potassium carbonate (1.4 g, 10 mmol) and potassium iodide (0.43 g, 2.6 mmol) were added and the stirred
15 mixture was heated at reflux temperature for 60 h. The reaction mixture was filtered, the filtrate concentrated in vacuo and the remainder was redissolved in diethyl ether (50 ml). The product was precipitated as the hydrochloride salt by dropwise addition of a 2.6 M solution of hydrogen chloride in diethyl ether (2.0 ml, 5.2 mmol). The precipitate was collected by filtration and dried in vacuo,
20 affording 1.3 g (63 %) of the 1-(3-(2,8-dibromo-10,11-dihydro-dibenz[b,f]-azepin-5-yl)-propyl)-4-piperidine-carboxylic acid ethyl ester hydrochloride as a powder, which was used in the next step without further purification.

The above ester hydrochloride (1.30 g, 2.2 mmol) was dissolved in ethanol (15
25 ml) and 2N sodium hydroxide (4.0 ml) was added. The solution was stirred for 1.5 h at room temperature, acidified by addition of 1 N hydrochloric acid to pH 1 and ethanol was evaporated in vacuo. The aqueous suspension was washed

56

with diethyl ether and filtered. Recrystallisation of the collected solid from ethanol afforded 0.25 g, (19 %) of the title compound.

M.p. 165 - 166 °C.

5 Calculated for $C_{23}H_{26}N_2Br_2O_2$, HCl, 0.5 H_2O , 0.5 C_2H_6O :


C, 48.79 %; H, 5.29 %; N, 4.74 %; Found:

C, 48.64 %; H, 5.39 %; N, 4.58 %.

10

EXAMPLE 22

1-(3-(3,7-Dichloro-10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-4-piperidinecarboxylic acid hydrochloride.



15

3,7-Dichloro-10,11-dihydro-5H-dibenz[b,f]azepine (4.6 g, 17 mmol) was dissolved in toluene (30 ml) and 3-chloropropionyl chloride (2.3 ml, 24 mmol) was added. The resulting mixture was heated at 90 °C for 3 h, stirred at room

temperature for 3 days and heated at 90 °C for 3 h. After cooling, ethyl acetate (150 ml) was added and the mixture was washed with water (2 x 100 ml). The organic phase was dried (MgSO_4) and evaporated in vacuo. The residue was purified by column chromatography on silica gel (600 ml), using a mixture of
5 ethyl acetate and heptane (1:4) as eluent. This afforded 4.7 g (76 %) 3-chloro-1-(3,7-dichloro-10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propanone as an oil.

TLC: R_f = 0.14 (SiO_2 : ethyl acetate/heptane = 1:4)

10 Lithium aluminum hydride (0.44 g, 11.6 mmol) was dissolved in dry tetrahydrofuran (15 ml) and the solution was cooled to 0 °C. Concentrated sulphuric acid (0.31 ml, 5.8 mmol) was slowly and cautiously added using a syringe. The suspension was stirred at room temperature for 0.5 h. A solution of the above amide (2.0 g, 5.8 mmol) in dry tetrahydrofuran (15 ml) was added
15 dropwise to the suspension. The resulting mixture was stirred at room temperature for 1 h. The reaction was quenched by addition of water (1.4 ml), 2 N sodium hydroxide (0.5 ml), water (4 ml), and potassium carbonate (5.0 g). The mixture was filtered, and the filter cake was washed with dry tetrahydrofuran. The combined filtrates were evaporated in vacuo to give 1.4 g (69 %) 5-(3-
20 chloropropyl)-3,7-dichloro-10,11-dihydro-5H-dibenz[b,f]azepine as an oil.

TLC: R_f = 0.66 (SiO_2 : ethyl acetate/heptane = 1:2)

The above chloride (1.3 g, 3.8 mmol) was dissolved in methyl ethyl ketone (20
25 ml). Potassium iodide (0.63 g, 3.8 mmol), potassium carbonate (3.2 g, 23 mmol), and 4-piperidinecarboxylic acid ethyl ester (1.2 ml, 7.6 mmol) were added, and the resulting mixture was heated at reflux temperature for 24 h.

After cooling, ethyl acetate (100 ml) was added and the mixture was washed with water (2 x 100 ml). The organic phase was dried (MgSO_4) and evaporated in vacuo. The residue was purified by column chromatography on silica gel (600 ml), using a mixture of ethyl acetate and heptane (1:2) as eluent. This
5 afforded 0.84 g (48 %) 1-(3-(3,7-dichloro-10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-4-piperidinecarboxylic acid ethyl ester as an oil.

TLC: R_f = 0.22 (SiO_2 : ethyl acetate/heptane = 1:2)

10 The above ethyl ester (0.8 g, 1.7 mmol) was dissolved in ethanol (20 ml), and water (10 ml) and 1 N sodium hydroxide (1.7 ml) were added. The resulting mixture was stirred at room temperature for 24 h. Water (100 ml) was added and the mixture was washed with diethyl ether (2 x 60 ml). The pH of the aqueous phase was adjusted to 1 with 5 N hydrochloric acid, and the aqueous
15 phase was extracted with dichloromethane (2 x 100 ml). The combined organic extracts were dried (MgSO_4) and evaporated in vacuo to give an oil. 2-Propanol (15 ml) was added, and the resulting precipitate was filtered off and washed with 2-propanol. Drying in vacuo at 50 °C for 24 h afforded 0.33 g (41 %) of the title compound.

20

M.p. 237 - 239 °C

MS(EI) (m/z): 432 (M^+ , 14 %), 303 (43 %), 142 (100 %).


^1H NMR (400 MHz, CDCl_3): δ_{H} 1.72 - 2.02 (m, 5H), 2.48 (m, 1H), 2.85 (m, 2H),
3.05 (m, 2H), 3.08 (s, 4H), 3.39 (m, 2H), 3.80 (t, 2H), 7.03 (dd, 2H), 7.17 (d,

25 2H), 7.22 (d, 2H).

EXAMPLE 23

1-(3-(3-Methyl-10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-1-propyl)-4-piperidinecarboxylic acid hydrochloride

5



-
- 10 5-(3-Bromo-1-propylidene)-3-methyl-10,11-dihydro-5H-dibenzo[a,d]cycloheptene (1.05 g, 3.3 mmol, prepared as described in WO 9518793) was dissolved in methyl ethyl ketone (20 ml). Potassium iodide (0.44 g, 6.7 mmol), potassium carbonate (2.76 g, 20 mmol), and 4-piperidinecarboxylic acid ethyl ester (0.77 ml, 5.0 mmol) were added and the resulting mixture was heated at reflux
- 15 temperature for 1 h. After cooling, ethyl acetate (60 ml) was added and the mixture was washed with water (2 x 60 ml). The organic phase was dried (MgSO₄) and evaporated *in vacuo*. The residue (1.7 g) was purified by column chromatography on silica gel (600 ml), using a mixture of ethyl acetate and heptane (1:2) as eluent. This afforded 0.97 g (73 %) 1-(3-(3-methyl-10,11-
- 20 dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-1-propyl)-4-piperidinecarboxylic acid ethyl ester as an oil.

TLC: $R_f = 0.20$ (SiO_2 : ethyl acetate/heptane = 1:2)

The above ethyl ester (0.91 g, 1.7 mmol) was dissolved in ethanol (15 ml), and water (5 ml) and 1 N sodium hydroxide (2.7 ml) were added. The resulting mixture was stirred at room temperature for 24 h. The reaction mixture was evaporated in vacuo. Water (100 ml) and diethyl ether (70 ml) were added, and the phases were separated. The pH of the aqueous phase was adjusted to 2 with 1 N hydrochloric acid, and the aqueous phase was extracted with dichloromethane (3 x 200 ml). The combined organic extracts were dried (MgSO₄) and evaporated in vacuo to give 0.12 g (13 %) of the title compound. More material separated from the acidic aqueous phase on standing. The precipitate was filtered off, washed with water, and dried to give 0.34 g (36 %) of the title compound.

15 M.p. >250 °C

Calculated for C₂₅H₂₉NO₂, HCl:

C, 72.89 %; H, 7.34 %; N, 3.40 %. Found:

C, 72.65 %; H, 7.43 %; N, 3.26%.


20

EXAMPLE 24

1-(3-(3,7-Dimethyl-10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-4-

25 piperidinecarboxylic acid

61



- To a solution of 3,7-dimethyl-10,11-dihydro-5H-dibenz[b,f]azepine (6.45 g,
5 0.029 mol; prepared similarly as described in Brit. Pat. 792615, 1958) and
3-bromo-1-propyl tetrahydro-2-pyranyl ether (8.3 g, 0.037 mol) in dry
benzene (80 ml) a suspension of sodium amide (3.2 g, 0.041 mol, 50%wt
suspension in toluene) was added. The reaction mixture was heated at
reflux temperature for 20 h, allowed to cool, and water (20 ml) was added.
- 10 The phases were separated, the solvent was evaporated from the organic
phase and the residue was dissolved in a mixture of methanol (100 ml) and
5 N HCl (30 ml). The mixture was then heated at reflux temperature for 15
minutes, methanol was evaporated and the mixture was extracted with
benzene (2 x 150 ml). The combined organic extracts were dried (K_2CO_3),
15 filtered and the solvent was evaporated under vacuum. The residue was
purified by chromatography on silica gel (50 g) using first benzene as eluent
to separate the starting material. Then, using chloroform as eluent, 2.4 g of
3-(3,7-dimethyl-10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propanol as an
oil was separated.

The above alcohol (2.4 g, 0.0087 mol) was dissolved in benzene (80 ml) and then triethyl amine (3.0 ml) was added. After addition of methane sulfonylchloride (1.3 g, 0.0114 mol) the reaction mixture was stirred for 2 h. Water was added and the phases were separated. The organic phase was
5 dried (MgSO_4) and the solvent was evaporated under vacuum. The residue was dissolved in acetone (50 ml). To this solution 4-piperidinecarboxylic acid ethyl ester (2.0 g, 0.0127 mol) and potassium carbonate (3.0 g, 0.0217 mol) were added, and the mixture was heated at reflux temperature for 24 h. The mixture was allowed to cool, then filtered and the solvent was
10 evaporated under vacuum to give a residue, which was further purified by chromatography on silica gel (40 g) using chloroform as eluent. This afforded 2.6 g of 1-(3-(3,7-dimethyl-10,11-dihydro-5H-dibenz[b,f]-azepin-5-yl)-1-propyl)-4-piperidinecarboxylic acid ethyl ester as an oil.

15 The above ester (2.4 g, 0.057 mol) was dissolved in ethanol (50 ml) and 5 N NaOH (3 ml) was added. The mixture was stirred at 40 °C for 16 h, and ethanol was evaporated under vacuum to give a residue, which was dissolved in water (20 ml). Acetic acid (3 ml) was added to the resulting solution, and the mixture was extracted with dichloromethane (50 ml). The
20 organic extract was dried (MgSO_4), and the solvent was evaporated under vacuum. Diethyl ether (50 ml) was added to the residue, affording after filtration and drying 1.85 g (82%) of the title compound as a solid.

M.p. 207 - 209 °C.


25 Calculated for $\text{C}_{25}\text{H}_{32}\text{N}_2\text{O}_2$, 0.25 H_2O :

C, 75.63 %; H, 8.25 %; N, 7.06 %; Found:

C, 75.58 %; H, 8.30 %, N, 6.89%.

EXAMPLE 25

1-(3-(3-Dimethylamino-10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-4-
5 piperidinecarboxylic acid



-
- 10 A suspension of sodium amide (2.6 g, 0.033 mol, 50%wt suspension in toluene) was added to a solution of 3-dimethylamino-10,11-dihydro-5H-dibenz[b,f]azepine (6.1 g, 0.0256 mol, prepared similarly as described in Brit. Pat., 1040739,1966) in dry benzene (60 ml). The reaction mixture was heated to 70 °C for 1 h. 3-Bromo-1-propyl-tetrahydro-2-pyranyl ether (7.35
- 15 g, 0.033 mol) was added and the mixture was heated at reflux temperature for 20 h. To the cooled reaction mixture, water (20 ml) was added, and the phases were separated. The solvent was evaporated from the organic phase and the residue was dissolved in a mixture of methanol (100 ml) and 5 N

HCl (30 ml). The mixture was heated at reflux temperature for 15 minutes, and methanol was evaporated. Water (50 ml) was added, pH was adjusted to 8-9 with aqueous ammonia and the mixture was extracted with benzene (2 x 150 ml). The combined organic extracts were dried (K_2CO_3), filtered
5 and the solvent was evaporated under vacuum. The residue, was purified by chromatography on silica gel (50 g) using first benzene as eluent to isolate the starting material. Then, using chloroform as eluent, 3.5 g of 3-(3-dimethylamino-10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propanol as an oil was isolated.

10

The above alcohol (3.5 g, 0.0118 mol) was dissolved in benzene (100 ml) and triethylamine (4.0 ml) and methane sulfonylchloride (1.79, 0.0148 mol) were added. The reaction mixture was stirred for 2 h. After addition of water, the phases were separated. The organic phase was dried ($MgSO_4$)
15 and the solvent was evaporated under vacuum to give a residue, which was dissolved in acetone (50 ml). To this solution 4-piperidinecarboxylic acid ethyl ester (2.8 g, 0.0178 mol) and potassium carbonate (4.13, 0.03 mol) were added and the mixture was heated at reflux temperature for 24 h. The mixture was allowed to cool, filtered and the solvent was evaporated under
20 vacuum to give a residue, which was further purified by chromatography on silica gel (50 g) using ethyl acetate as eluent. This afforded 3.1 g of 1-(3-(3-dimethylamino-10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-4-piperidinecarboxylic acid ethyl ester as an oil.

25 The above ester (1.95 g, 0.0045 mmol) was dissolved in ethanol (40 ml) and 5 N NaOH (3 ml) was added. The mixture was stirred at 40 °C for 8 h, and ethanol was evaporated under vacuum. The residue was dissolved in

65

water (20 ml). Acetic acid (3 ml) was added to the resulting solution and the mixture was extracted with dichloromethane (50 ml). The organic extract was dried (MgSO_4), and the solvent was evaporated under vacuum. Diethyl ether (50 ml) was added to the residue, affording 1.67 g (91 %) of
5 the title compound as a solid after filtration and drying.

M.p. 198-202°C.


Calculated for $\text{C}_{25}\text{H}_{33}\text{N}_3\text{O}_2$, 0.25 H_2O :

C, 72.87 %; H, 8.19 %; N, 10.20 %, Found:

10 C, 72.73 %; H, 8.32 %; N, 10.00 %.

EXAMPLE 26

(R)-1-(3-(10,11-Dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-1-propyl)-2-
15 piperidinecarboxylic acid hydrochloride.



2-Piperidinecarboxylic acid (26 g, 0.201 mol) and (+)-tartaric acid (31.2 g, 0.208 mol) were suspended in a mixture of ethanol (400 ml) and water (25 ml). The mixture was heated to 80 °C, and the solution was allowed to cool to room temperature. The precipitated solid was filtered off, washed with ethanol and dried to give 25.4 g (45 %) (R)-(+)-2-piperidinecarboxylic acid (+)-tartrate. The mother liquor was evaporated in vacuo, and the residue was dissolved in water (120 ml). A solution of potassium hydroxide (6.5 g, 116 mmol) in water (13 ml) was added, and the precipitated mono potassium tartrate was removed by filtration. Evaporation of the filtrate in vacuo afforded 19.5 g crude (S)-(-)-2-piperidinecarboxylic acid.

The above (+)-tartrate (20 g, 72 mmol) was dissolved in water (60 ml) and a solution of potassium hydroxide (4.0 g, 72 mmol) in water (8 ml) was added. The precipitated mono potassium tartrate was removed by filtration and washed with water. The mother liquor was evaporated in vacuo, and the residue was suspended in ethanol (100 ml) and evaporated in vacuo. Suspension in ethanol and evaporation in vacuo was repeated twice. The residue was suspended in ethanol (200 ml), and thionyl chloride (24 ml, 0.28 mol) was added dropwise. The resulting mixture was heated at 70 °C for 2 h, cooled to room temperature and evaporated in vacuo. Diethyl ether (40 ml) and ethanol (1 ml) were added and the suspension was stirred for 30 minutes. The solid was filtered off, washed with diethyl ether and dried to give 11.4 g (82 %) of (R)-(+)-2-piperidinecarboxylic acid ethyl ester hydrochloride.

$[\alpha]^{25}_D = +10.5^\circ$ (c = 4.5 % in water)

Gas Chromatography of N-acetyl derivative: $R_t = 46.4$ minutes. Enantiomeric excess = 97.9 %

- 5 (GC was run on a Chrompac CP 9000 Gas Chromatograph with F.I.D. detection using a Chrompac CP-Cyclodextrin capillary column, 25 m long, 0.25 mm internal diameter, using a split flow of 40 ml/min, 180 kPa, a linear gas velocity of 27.6 cm/s, inlet and detector temperatures of 200 °C and a column temperature of 120 °C).

10

- 5-(3-Bromo-1-propylidene)-10,11-dihydro-5H-dibenzo[a,d]cycloheptene (2.7 g, 8.6 mmol, prepared as described in WO 9518793), potassium carbonate (7.14 g, 52 mmol), potassium iodide (1.4 g 8.6 mmol), and (R)-(+)-2-piperidine-carboxylic acid ethyl ester hydrochloride (3.3 g, 17 mmol) were
15 mixed in methyl ethyl ketone (50 ml) and heated at reflux temperature for 3 days. After cooling to room temperature, ethyl acetate (100 ml) was added and the mixture was washed with water (2 x 100 ml). The organic phase was dried ($MgSO_4$) and evaporated in vacuo. The residue was purified by column chromatography on silica gel (600 ml), using a mixture of ethyl
20 acetate and heptane (1:4) as eluent. This afforded 2.62 g (78 %) of (R)-1-(3-(10,11-dihydro-5H-dibenzo[a,d]cyclo-hepten-5-ylidene)-1-propyl)-2-piperidinecarboxylic acid ethyl ester as an oil.

TLC: $R_f = 0.28$ (SiO_2 : ethyl acetate/heptane = 1:4)

25

The above ester (2.6 g, 6.7 mmol) was dissolved in a mixture of ethanol (25 ml) and 1,4-dioxane (2 ml). 1 N Sodium hydroxide (6.7 ml) was added and

the mixture was stirred at room temperature for 16 h. 1 N Sodium hydroxide (1.3 ml) was added and the mixture was stirred at room temperature for 4 h. 1 N Sodium hydroxide (6.9 ml) and ethanol (10 ml) were added and the mixture was stirred at room temperature for 16 h. 1 N Sodium hydroxide (6.9 ml) was added and the mixture was stirred at room temperature for 16 h. 1 N Sodium hydroxide (6.7 ml) was added and the mixture was stirred at room temperature for 3 days. Water (100 ml) was added and the mixture was washed with diethyl ether (3 x 100 ml). The pH of the aqueous phase was adjusted to 1 with 5 N hydrochloric acid, and the aqueous phase was extracted with dichloromethane (150 ml). The organic extract was dried (MgSO₄) and evaporated in vacuo to give an oil. Acetone (50 ml) was added and the solution was evaporated in vacuo. The acetone treatment was repeated. The solid was suspended in acetone and filtered off. Drying in vacuo at 50 °C for 24 h afforded 1.2 g (51 %) of the title compound as an amorphous powder.

MS(EI) (m/z): 316 (M⁺, 1 %), 142 (100 %).


¹H NMR (400 MHz, CDCl₃): δ_H 1.45 - 1.75 (m, 5H), 2.1 (m, 1H), 2.4-2.55 (m, 2H), 2.75-3.4 (m, 8H), 3.90 (bs, 1H), 5.80 (t, 1H), 7.05-7.23 (m, 8H).

20

EXAMPLE 27

25 (S)-1-(3-(10,11-Dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-1-propyl)-2-piperidinecarboxylic acid hydrochloride.

69



-
- Crude (S)-(-)-2-piperidinecarboxylic acid (19.5 g, prepared as described in
- 5 example 26) was suspended in ethanol (250 ml), and thionyl chloride (40 ml, 0.46 mol) was added dropwise. After addition was complete, the suspension was heated at reflux temperature for 2 h. The mixture was filtered hot, and the filtrate was cooled to room temperature. Filtration and evaporation in vacuo afforded an oil, which was crystallised by rubbing.
- 10 Ethanol (10 ml) was added, followed by slow addition of diethyl ether (150 ml). The precipitated solid was filtered off, washed with diethyl ether and dried by suction to give 13.8 g (35 % calculated from 2-piperidinecarboxylic acid) (S)-(-)-2-piperidinecarboxylic acid ethyl ester hydrochloride.
- 15 $[\alpha]^{25}_D = -10.7^\circ$ (c = 4.5 % in water)
- Gas Chromatography (run as described in example 26) of N-acetyl derivative: $R_t = 47.2$ minutes. Enantiomeric excess = 96 %.

5-(3-Bromo-1-propylidene)-10,11-dihydro-5H-dibenzo[a,d]cycloheptene (2.7

- g, 8.6 mmol, prepared as described in WO 9518793), potassium carbonate (7.14 g, 52 mmol), potassium iodide (1.4 g, 8.6 mmol), and (S)-(-)-2-piperidinecarboxylic acid ethyl ester hydrochloride (3.3 g, 17 mmol) were mixed in methyl ethyl ketone (50 ml) and heated at reflux temperature for 3 days.
- 5 After cooling to room temperature, ethyl acetate (100 ml) was added and the mixture was washed with water (2 x 100 ml), dried (MgSO_4) and evaporated in vacuo. The residue was purified by column chromatography on silica gel (600 ml), using a mixture of ethyl acetate and heptane (1:4) as eluent. This afforded 2.3 g (69 %) of (S)-1-(3-(10,11-dihydro-5H-dibenzo-
- 10 [a,d]cyclohepten-5-ylidene)-1-propyl)-2-piperidinecarboxylic acid ethyl ester as an oil.

TLC: R_f = 0.22 (SiO_2 : ethyl acetate/heptane = 1:4)

- 15 The above ester (2.3 g, 5.9 mmol) was dissolved in a mixture of ethanol (25 ml) and 1,4-dioxane (2 ml). 1 N Sodium hydroxide (5.9 ml) was added and the mixture was stirred at room temperature for 16 h. 1 N Sodium hydroxide (1.18 ml) was added and the mixture was stirred at room temperature for 4 h. 1 N Sodium hydroxide (5.9 ml) was added and the mixture was
- 20 stirred at room temperature for 16 h. 1 N Sodium hydroxide (5.9 ml) was added and the mixture was stirred at room temperature for 16 h. 1 N Sodium hydroxide (5.9 ml) was added and the mixture was stirred at room temperature for 3 days. Water (100 ml) was added and the mixture was washed with diethyl ether (3 x 100 ml). The pH of the aqueous phase was
- 25 adjusted to 1 with 5 N hydrochloric acid, and the aqueous phase was extracted with dichloromethane (150 ml). The organic extract was dried (MgSO_4) and evaporated in vacuo to give an oil. Acetone (50 ml) was added

71

and the solution was evaporated in vacuo. The acetone treatment was repeated again. The solid was suspended in acetone, filtered off and dried in vacuo at 50 °C for 24 h to give 0.79 g (37 %) of the title compound as an amorphous powder.

5

MS(EI) (m/z): 362 (M⁺, 1 %), 142 (100 %).


¹H NMR (400 MHz, CDCl₃): δ_H 1.45 - 1.75 (m, 5H), 2.1 (m, 1H), 2.4-2.55 (m, 2H), 2.75-3.4 (m, 8H), 3.80 (bs, 1H), 5.80 (t, 1H), 7.05-7.23 (m, 8H).

10

- 72 -

CLAIMS


1. A compound of formula I




10 wherein R^1 and R^2 independently are hydrogen, halogen, trifluoromethyl, NR^6R^7 , hydroxy, C_{1-6} -alkyl or C_{1-6} -alkoxy; and
 Y is $>\underline{N}-CH_2-$, $>\underline{CH}-CH_2-$ or $>\underline{C}=CH-$ wherein only the underscored atom participates in the ring system; and

15 X is $-O-$, $-S-$, $-C(R^6R^7)-$, $-CH_2CH_2-$, $-CH=CH-CH_2-$, $-CH_2-CH=CH-$, $-CH_2-$, $(C=O)-$, $-(C=O)-CH_2-$, $-CH_2CH_2CH_2-$, $-CH=CH-$, $-N(R^8)-(C=O)-$, $-(C=O)-N(R^8)-$, $-O-CH_2-$, $-CH_2-O-$, $-S-CH_2-$, $-CH_2-S-$, $-(C=O)-$, $-N(R^9)-$ or $-(S=O)-$ wherein R^6 , R^7 , R^8 and R^9 independently are hydrogen or C_{1-6} -alkyl; and
 r is 1, 2 or 3; and
 Z is selected from


20



25



30



- 73 -

wherein n is 1 or 2; and

R³ is -(CH₂)_mOH or -(CH₂)_pCOR⁴ wherein m is 0, 1, 2, 3, 4, 5 or 6 and p is 0 or 1 and wherein R⁴ is -OH, -NH₂, -NHOH or C₁₋₆-alkoxy; and

R⁵ is hydrogen, halogen, trifluoromethyl, hydroxy, C₁₋₆-alkyl or C₁₋₆-alkoxy; and

R¹⁰ is hydrogen, C₁₋₆-alkyl, C₁₋₆-alkoxy or phenyl optionally substituted with halogen, trifluoromethyl, hydroxy, C₁₋₆-alkyl or C₁₋₆-alkoxy; and

R¹¹ is hydrogen or C₁₋₆-alkyl; and

... is optionally a single bond or a double bond;

or a pharmaceutically acceptable salt thereof.

2. A compound according to claim 1 wherein R¹ and R² independently are hydrogen, halogen, N(CH₃)₂ or methyl.

3. A compound according to any one of the preceding claims wherein Y is >N-CH₂- or >C=CH-.


4. A compound according to any one of the preceding claims wherein X represents -O-, -S-, -CH₂CH₂-, -CH₂CH₂CH₂-, -CH=CH-, -O-CH₂-, -CH₂-O-, -S-CH₂-, -CH₂-S-, -(C=O)- or -(S=O)-.

5. A compound according to the previous claim wherein X is -O- or -CH₂CH₂-.

6. A compound according to any one of the preceding claims wherein r is 2.

7. A compound according to any one of the preceding claims wherein Z represents

- 74 -



wherein n is 1

8. A compound according to the previous claim wherein R³ is $-(CH_2)OH$, $-CH_2CH_2OH$, $-COR^4$ or $-CH_2COR^4$ wherein R⁴ is OH or NH₂.

15

9. A compound according to claim 7 wherein R⁵ is hydrogen or C₁₋₆-alkyl.

10. A compound according to claim 7 wherein R¹⁰ is hydrogen or methyl, or phenyl substituted with halogen.

20

11. A compound according to claim 7 wherein R¹¹ is hydrogen or methyl.

12. A compound according to claim 7 wherein ... is a single bond.

25

13. A compound according to any of the claims 1 through 12 selected from the following:

1-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-3-piperidine-carboxamide;

30

- 75 -

- 1-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-4-piperidinecarboxylic acid;
- 5 1-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-2-piperidinecarboxylic acid;
- (1-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-3-piperidinyl)-methanol;
- 10 4-(4-Chlorophenyl)-1-(3-(10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-4-piperidinol;
- 4-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-2-piperazinecarboxylic acid;
- 15 (2S,4R)-1-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-4-hydroxy-2-pyrrolidinecarboxylic acid;
- 4-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-2-morpholinecarboxylic acid;
- 20 1-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-2-aziridinecarboxylic acid;
- 25 2-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-1,2,3,4-tetrahydro-4-isoquinolinecarboxylic acid;
- 1-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-4-methyl-[1,4]-diazepane-6-carboxylic acid;
- 30 2-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-1,2,3,4-tetra-

- 76 -

hydro-3-isoquinolinecarboxylic acid;

1-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-3-piperidinecarboxylic acid hydroxamide;

5

(4-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)piperazin-1-yl)acetic acid;

10

1-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-4-piperidinecarboxylic acid;

4-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-2-piperazinecarboxylic acid;

15

1-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-4-piperidineacetic acid;

1-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-4-piperidineacetic acid;

20

1-(3-(10,11-Dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-1-propyl)-4-piperidinecarboxylic acid;

25

(R)-1-(3-(10,11-Dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-1-propyl)-3-piperidinecarboxamide;

(R)-1-(3-(10,11-Dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-1-propyl)-2-pyrrolidinecarboxylic acid;

30

(S)-1-(3-(10,11-Dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-1-propyl)-2-pyrrolidinecarboxylic acid;

- 77 -

- 1-(3-(10,11-Dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-1-propyl)-2-piperidinecarboxylic acid;
- 5 1-(3-(10H-Phenoxazin-10-yl)-1-propyl)-4-piperidinecarboxylic acid;
- 1-(3-(3-Chloro-10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-4-piperidinecarboxylic acid;
- 10 1-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-3-piperidine-acetic acid;
- 15 1-(3-(10,11-Dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-3-quinuclidiniumcarboxylate;
- 1-(3-(2,8-Dibromo-10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-4-piperidinecarboxylic acid;
- 20 1-(3-(3,7-Dichloro-10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-4-piperidinecarboxylic acid;
- 25 1-(3-(3-Methyl-10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-1-propyl)-4-piperidinecarboxylic acid;
- 1-(3-(3,7-Dimethyl-10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-4-piperidinecarboxylic acid;
- 30 1-(3-(3-Dimethylamino-10,11-dihydro-5H-dibenz[b,f]azepin-5-yl)-1-propyl)-4-piperidinecarboxylic acid;

- 78 -

(R)-1-(3-(10,11-Dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-1-propyl)-
2-piperidinecarboxylic acid;

(S)-1-(3-(10,11-Dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-1-propyl)-
5 2-piperidinecarboxylic acid;

or a pharmaceutically acceptable salt thereof.

10 14. The use of a compound according to any of the claims 1 through 13
as a medicament.

15 15. The use of a compound according to any of the claims 1 through 13
for preparing a medicament for the treatment of neurogenic inflam-
mation.

16. The use of a compound according to any of the claims 1 through 13
for preparing a medicament for the treatment of insulin resistance in
NIDDM or aging.

20 17. A pharmaceutical composition comprising as active component a
compound according to any of the claims 1 through 13 together with a
pharmaceutically carrier or diluent.

25 18. A pharmaceutical composition suitable for treating neurogenic
inflammation comprising an effective amount of a compound according
to any one of the claims 1 through 13 together with a pharmaceutically
acceptable carrier or diluent.

30 19. A pharmaceutical composition suitable for treating insulin resistance
in NIDDM or aging comprising an effective amount of a compound
according to any one of the claims 1 through 13 together with a pharma-

- 79 -

ceutically acceptable carrier or diluent.

5 **20.** The pharmaceutical composition according to claims 17, 18 or 19 comprising between 0.5 mg and 1000 mg of the compound according to any one of the claims 1 through 13 per unit dose.

10 **21.** A method of treating neurogenic inflammation in a subject in need of such treatment comprising administering to said subject an effective amount of a compound according to any one of the claims 1 through 13.

15 **22.** A method of treating insulin resistance in NIDDM or aging in a subject in need of such treatment comprising administering to said subject an effective amount of a compound according to any one of the claims 1 through 13.


20 **23.** A method of treating neurogenic inflammation in a subject in need of such treatment comprising administering to said subject a pharmaceutical composition according to claim 18.

25 **24.** A method of treating insulin resistance in NIDDM or aging in a subject in need of such treatment comprising administering to said subject a pharmaceutical composition according to claim 19.

30 **25.** A method of preparing a compound according to claim 1,
CHARACTERIZED in

a) reacting a compound of formula II

- 80 -



(II)

5

wherein R¹, R², X, Y and r are as defined above and W is a suitable leaving group such as halogen, p-toluene sulphonate or mesylate, with a compound of formula III

10

HZ

(III)

wherein Z is as defined above to form a compound of formula I; or

15

b) hydrolyzing a compound of formula I, wherein R⁴ is C₁₋₆-alkoxy, to form a compound of formula I wherein R⁴ is OH.

26. Any novel feature or combination of features described herein.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 96/00139

A. CLASSIFICATION OF SUBJECT MATTER

IPC6: C07D 401/06, C07D 403/06, C07D 413/06, C07D 211/60, A61K 31/55, A61K 31/445
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS-ONLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GB 830709 A (G.D. SEARLE & CO.), 16 March 1960 (16.03.60) --	1-4,6-8,10, 12,17-20,25
X	Chemical Abstracts, Volume 76, No 23, 5 June 1972, (05.06.72), (Columbus, Ohio, USA), page 460, THE ABSTRACT No 140577n, JP, 7204069, A, (Nakanishi, Michio et al) 4 February 1972 (04.02.72) --	1-4,6-7, 17-20,25
X	Chemical Abstracts, Volume 81, No 5, 5 August 1974, (05.08.74), (Columbus, Ohio, USA), page 424, THE ABSTRACT No 25566z, JP, 7330064, A, (Nakanishi, Michio et al) 23 December 1968 (23.12.68) --	1-7,17-20,25



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

& document member of the same patent family

Date of the actual completion of the international search

3 July 1996

Date of mailing of the international search report

06-08-1996

Name and mailing address of the ISA/

Swedish Patent Office
Box 5055, S-102 42 STOCKHOLM
Facsimile No. +46 8 666 02 86

Authorized officer

Göran Karlsson
Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 96/00139

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Collect.Czech.Chem.Comm., Volume 59, 1994, Karel Sindélar et al, "Antihistamine, Substances. Tricyclic Analogues of N-(4,4-Dipheny, -3-Butene-1-YL,)Nipecotic Acid and some Related Compounds", page 667 - page 675, see compounds I-III --	1-8,12, 17-20,25
X	US 3326896 A (ELVIN A. HOLSTIUS ET AL), 20 June 1967 (20.06.67) --	1-4,6-8, 17-20,25
X	FR 1296519 A (ABBOTT LABORATORIES), 14 May 1962 (14.05.62) --	1-4,6-8, 17-20,25
X	US 4139632 A (NORIYASU HIROSE ET AL), 13 February 1979 (13.02.79) --	1,3-4,6-7, 17-20,25
E,X	WO 9518793 A1 (NOVO NORDISK A/S), 13 July 1995 (13.07.95) -- -----	1-8,12, 17-20,25

INTERNATIONAL SEARCH REPORT

Int. application No.

PCT/DK 96/00139

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 14, 21-24
because they relate to subject matter not required to be searched by this Authority, namely:
A method for treatment of the human or animal body by therapy,
see rule 39.1
2. ☒ Claims Nos.: 26
because they relate to parts of the international application that do not comply with the prescribed requirements to such
an extent that no meaningful international search can be carried out, specifically:
Claim 26 does not clearly define the matter for which protection is
sought, see Article 6
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

5776.6

INTERNATIONAL SEARCH REPORT
Information on patent family members

01/04/96

International application No.

PCT/DK 96/00139

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB-A- 830709	16/03/60	BE-A- 562299	00/00/00
		CH-A- 366043	00/00/00
		DE-B- 1089386	00/00/00
		FR-A- 1231657	00/00/00
		NL-C- 99690	00/00/00
US-A- 3326896	20/06/67	BE-A- 658764	26/07/65
		BR-D- 6566386	00/00/00
		CH-A- 454871	00/00/00
		DE-A- 1545882	19/03/70
		FR-A- 1426451	00/00/00
		GB-A- 1051221	00/00/00
		NL-A- 6500845	26/07/65
		SE-B- 305449	28/10/68
FR-A- 1296519	14/05/62	NONE	
US-A- 4139632	13/02/79	DE-A, A- 2636866	03/03/77
		FR-A, B- 2321894	25/03/77
		GB-A- 1517385	12/07/78
		JP-C- 1096522	14/05/82
		JP-A- 52027779	02/03/77
		JP-B- 56037990	03/09/81
		NL-A- 7609458	02/03/77
		SE-B, C- 412233	25/02/80
		SE-A- 7609466	01/03/77
WO-A1- 9518793	13/07/95	NONE	